

541, 460

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
26 August 2004 (26.08.2004)

PCT

(10) International Publication Number  
**WO 2004/072083 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 495/04**,  
487/04, 471/14, 471/04, 498/04, A61K 31/5517, A61P  
15/00

(21) International Application Number:  
PCT/EP2004/001304

(22) International Filing Date: 12 February 2004 (12.02.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
03003394.8 14 February 2003 (14.02.2003) EP

(71) Applicant (for all designated States except US): **FER-  
RING B.V.** [NL/NL]; P.O. Box 3129 2130 KC, Polarisav-  
enue 144, NL-2132 JX Hoofddorp (NL).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HUDSON, Peter**  
[GB/DK]; Amagerbrogade 64A, 2nd Floor, DK-2300  
Copenhagen S. (DK). **BATT, Andrzej, Roman** [GB/GB];  
41 Kent Road, Southampton SO17 2LJ (GB). **HEENEY,**  
**Celine, Marguerite, Simone** [FR/GB]; 59 Cedar Road,  
Portsmouth, Southampton SO14 6TQ (GB). **BAXTER,**

**Andrew, John** [GB/GB]; 15 Stapleford Close, Woodley,  
Romsey, Hants SO51 7HU (GB). **ROE, Michael, Bryan**  
[GB/GB]; 1B Cambridge Road, Southampton SO14 6RB  
(GB). **ROBSON, Peter, Andrew** [GB/GB]; 5 Eldridge  
Gardens, Romsey, Hampshire SO51 7RT (GB).


(74) Agent: **AWAPATENT AB**; Box 5117, S-200 71 Malmö  
(SE).

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,  
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,  
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,  
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,  
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,  
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,  
ZW.

(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Euro-  
pean (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,  
GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK,  
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,  
ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: **BENZAMIDE DERIVATIVES AS OXYTOCIN AGONISTS AND VASOPRESSIN ANTAGONISTS**



(57) Abstract: Novel compounds according to general formula 1, wherein G<sup>1</sup> is NR<sup>5</sup>R<sup>6</sup> or a fused polycyclic group that are specific OT receptor agonists and/or V<sub>1a</sub> receptor antagonists. Pharmaceutical compositions comprising such compounds are useful in the treatment of, *inter alia*, primary dysmenorrhoea.

WO 2004/072083 A1



**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

## BENZAMIDE DERIVATIVES AS OXYTOCIN AGONISTS AND VASOPRESSIN ANTAGONISTS


## FIELD OF THE INVENTION

The present invention relates to novel non-peptide oxytocin agonists and to pharmaceutical compositions comprising such compounds. The present invention also relates to the use of non-peptide oxytocin agonists for the treatment of certain physiological disorders, such as erectile dysfunction and primary dysmenorrhoea.

## BACKGROUND

### Neurophyseal hormones

10 The neurophyseal hormones oxytocin (OT) and vasopressin (VP) are cyclic nonapeptides secreted by the posterior pituitary gland. The structure of oxytocin is shown below.



Oxytocin - *cyclo*<sup>1,6</sup>-Cys-Tyr-Ile-Gln-Asn-Cys-Pro-Leu-Gly-NH<sub>2</sub>

15

Vasopressin differs from oxytocin in that it has phenylalanine at position 3 in place of isoleucine and arginine at position 8 in place of leucine. Both hormones are synthesised *in vivo* as larger precursors, neurophysins, which are subject to post-translational processing to release the mature peptides. OT and VP act through a family of heptahelical receptors. Only one OT receptor has so far been well characterised, while three VP receptors are known. These are designated the  $V_{1a}$ ,  $V_{1b}$  and  $V_2$  receptors.

The first target organs to be identified for OT were the uterus, where it is implicated in the onset and progress of labour, and mammary glands, where it is involved in the regulation of milk expression. Other organs also express OT receptors, and it is clear that OT has a range of physiological roles that have not been fully elaborated yet. In particular, it has been suggested that OT acting in the CNS is involved in the erectile response in males, and in the regulation of female sexual arousal. For example, OT is erectogenic when administered *i.c.v.* to male rats. It also has erectogenic activity when given *i.v.*, but the doses required are up to two orders of magnitude greater, which is consistent with a central mode of action.

Vasopressin acts on the blood vessels, where it is a potent vasoconstrictor, and on the kidneys, where it promotes water reuptake leading to an antidiuretic effect.

#### 15 Oxytocin agonists and antagonists

A number of peptide analogues of OT are known in the literature. These include both agonists and antagonists. OT and its agonists are used, for example, to accelerate labour and to increase uterine muscle tone to control post-partum bleeding, and one antagonist, atosiban, has recently been registered as a treatment for pre-term labour. However, the peptidic nature of these compounds means that they are not likely to be bioavailable after oral dosing or to cross efficiently into the CNS. In order to get drugs that can be given orally and to be able to exploit the central effects of OT, attention has increasingly turned to non-peptides. As a result, there are many publications describing non-peptide OT antagonists in early-stage development. So far, however, there have been no reports of non-peptide OT agonists. This is not unexpected, as it is generally held that it is easier to find a receptor antagonist than an agonist.

So there remains a need for non-peptide OT receptor agonists. Such compounds should preferably be selective for the OT receptor over the VP receptors. They could be expected to show therapeutic utility in male and female sexual dysfunction, particularly male erectile dysfunction, in promoting labour, in controlling post-partum bleeding, in increasing milk let-down as well as a number of other indications.



Vasopressin agonists and antagonists

The  $V_{1a}$  receptor is normally acted upon by the endogenous agonist ligand, arginine vasopressin (AVP). AVP also acts on the  $V_{1b}$  and  $V_2$  receptors. It exerts a variety of biological effects in mammals including regulation of water and solute excretion by the kidney. AVP is structurally related to OT.

The  $V_{1a}$ ,  $V_{1b}$ , and  $V_2$ , as well as the OT receptors, are members of the super-family of seven transmembrane receptors known as G-protein coupled receptors. The  $V_{1a}$  receptor mediates phospholipase C activation and intracellular calcium mobilisation. Localisation of the receptors includes blood platelets, blood vessels, hepatocytes, brain and uterus-cervix. Thus a  $V_{1a}$  antagonist may have effects on any or all of these tissues. For example, selective  $V_{1a}$  antagonists have been cited as having clinical utility in primary dysmenorrhoea, pre-term labour, hypertension, Raynaud's disease, brain oedema, motion sickness, small cell lung cancer, depression, anxiety, hyponatremia, liver cirrhosis and congestive heart failure.

With respect to primary dysmenorrhoea it has been proposed that myometrial activity is markedly increased in women with primary dysmenorrhoea during menstruation. It is proposed that the myometrial ischemia caused by increased uterine contractility might explain the menstrual pain. Furthermore, on the first day of menstruation, higher plasma concentrations of vasopressin have been measured in dysmenorrhoeal women than in controls.

In healthy women without dysmenorrhoea, intravenous infusion of lysine-vasopressin resulted in decreased uterine blood flow, increased uterine contractility and slight to moderate like-like pain, these effects being inhibited by a selective human  $V_{1a}$  receptor antagonist. (Bossmar T; Brouard R; Doberl A; Akerlund M, Department of Obstetrics and Gynaecology, University Hospital of Lund, Sweden BRITISH JOURNAL OF OBSTETRICS AND GYNAECOLOGY (1997 Apr), 104(4), 471-7.). Also, it is known that vasopressin contracts human uterine arteries in a dose-dependent and  $V_{1a}$ -mediated fashion.


The above evidence suggests that a  $V_{1a}$  antagonist would be an appropriate and effective treatment for primary dysmenorrhoea. Further evidence is taken from the clinical study carried out on the selective  $V_{1a}$  antagonist

SR49059 ("Effect of SR49059, an orally active  $V_{1a}$  vasopressin receptor antagonist, in the prevention of dysmenorrhea". Brouard, R.; Bossmar, T.; Fournie-Lloret, D.; Chassard, D.; Akerlund, M. Sanofi Recherche, Clinical Development, Paris, Fr. BJOG (2000), 107(5), 614-619.). It was found that

there was a dose-related decrease in pain and a dose-related decrease in the amount of additional pain-killer taken compared to patients taking placebo.


## SUMMARY OF THE INVENTION

- 5 The present invention is based on a series of potent and specific OT receptor agonists (herein also called OT agonists) and/or VP receptor antagonists (herein also called VP antagonists) according to general formula 1:



1

wherein  $G^1$  is a group according to general formula 2, 3, 4, 5, 6 or 7:



wherein:

- 10 -  $A^1$  is  $CH_2$ ,  $CH(OH)$ ,  $NH$ ,  $N$ -alkyl,  $O$  or  $S$ ;  
 -  $A^2$  is  $CH_2$ ,  $CH(OH)$ ,  $C(=O)$  or  $NH$ ;  
 -  $A^3$  is  $S$ ,  $NH$ ,  $N$ -alkyl,  $-CH=CH-$  or  $-CH=N-$ ;  
 -  $A^4$  and  $A^5$  are each  $CH$  or  $N$ ;  
 -  $A^6$  is  $CH_2$ ,  $NH$ ,  $N$ -alkyl or  $O$ ;  
 15 -  $A^7$  and  $A^{11}$  are  $C$  or  $N$ ;  
 -  $A^8$  and  $A^9$  are  $CH$ ,  $N$ ,  $NH$ ,  $N(CH_2)_dR^7$  or  $S$ ;

- $A^{10}$  is  $-\text{CH}=\text{CH}-$ ,  $\text{CH}$ ,  $\text{N}$ ,  $\text{NH}$ ,  $\text{N}-(\text{CH}_2)_d-\text{R}^7$  or  $\text{S}$ ;
  - $A^{12}$  and  $A^{13}$  are  $\text{N}$  or  $\text{C}$  and
  - $A^{14}$ ,  $A^{15}$  and  $A^{16}$  are  $\text{NH}$ ,  $\text{N}-\text{CH}_3$ ,  $\text{S}$ ,  $\text{N}$  or  $\text{CH}$ ,
- provided that not more than one of  $A^8$ ,  $A^9$  and  $A^{10}$  is  $\text{NH}$ ,  $\text{N}-(\text{CH}_2)_d-\text{R}^7$  or  $\text{S}$ ;
- 5 that  $A^7$  and  $A^{11}$  are not both simultaneously  $\text{N}$ ; that neither  $A^7$  nor  $A^{11}$  is  $\text{N}$  if one of  $A^8$ ,  $A^9$  and  $A^{10}$  is  $\text{NH}$ ,  $\text{N}-(\text{CH}_2)_d-\text{R}^7$  or  $\text{S}$ ; that if  $A^{10}$  is not  $-\text{CH}=\text{CH}-$  then one of  $A^8$ ,  $A^9$  and  $A^{10}$  is  $\text{NH}$ ,  $\text{N}-(\text{CH}_2)_d-\text{R}^7$  or  $\text{S}$  or one of  $A^7$  and  $A^{11}$  is  $\text{N}$ ; that not more than one of  $A^{14}$ ,  $A^{15}$  and  $A^{16}$  is  $\text{NH}$ ,  $\text{N}-\text{CH}_3$  or  $\text{S}$ ; that  $A^{12}$  and  $A^{13}$  are not both simultaneously  $\text{N}$ ; that if one of  $A^{14}$ ,  $A^{15}$  and  $A^{16}$  is  $\text{NH}$ ,
- 10  $\text{N}-\text{CH}_3$  or  $\text{S}$  then  $A^{12}$  and  $A^{13}$  are both  $\text{C}$ ; and that one of  $A^{14}$ ,  $A^{15}$  and  $A^{16}$  is  $\text{NH}$ ,  $\text{N}-\text{CH}_3$  or  $\text{S}$  or one of  $A^{12}$  and  $A^{13}$  is  $\text{N}$ ;
- $\text{X}^1$  is  $\text{O}$  or  $\text{NH}$ ;
  - $\text{R}^1$ ,  $\text{R}^2$  and  $\text{R}^3$  are each  $\text{H}$ , alkyl,  $\text{O}$ -alkyl,  $\text{F}$ ,  $\text{Cl}$  or  $\text{Br}$ ;
  - $\text{R}^4$  is  $\text{H}$ , alkyl, optionally substituted phenyl, pyridyl, thienyl or furyl, or is
- 15  $-(\text{CH}_2)_e-\text{R}^8$ ;
- $\text{R}^5$  and  $\text{R}^6$  are each, independently of each other, alkyl,  $\text{Ar}$  or  $-(\text{CH}_2)_f-\text{Ar}$ , where  $\text{Ar}$  is optionally substituted phenyl or thienyl;
  - $\text{R}^7$  and  $\text{R}^8$  are each, independently of each other,  $\text{H}$ , alkyl, optionally substituted phenyl, pyridyl, thienyl or furyl,  $\text{F}$ ,  $\text{OH}$ ,  $\text{O}$ -alkyl,  $\text{S}$ -alkyl,  $\text{O}$ -acyl,  $\text{NH}_2$ ,
- 20  $\text{NH}$ -alkyl,  $\text{N}(\text{alkyl})_2$ ,  $\text{NH}$ -acyl,  $\text{N}(\text{alkyl})$ -acyl,  $\text{CO}_2\text{H}$ ,  $\text{CO}_2$ -alkyl,  $\text{CONH}_2$ ,  $\text{CONH}$ -alkyl,  $\text{CON}(\text{alkyl})_2$ ,  $\text{CN}$  or  $\text{CF}_3$ ;
- $a$  is 1 or 2,  $b$  is 1, 2 or 3,  $c$  is 1 or 2,  $d$  is 1, 2 or 3;  $e$  is 1, 2 or 3 and  $f$  is 1, 2 or 3.

According to a first aspect, the present invention relates to novel OT  
25 agonists and/or VP antagonists, and in particular specific antagonists of the  $\text{V}_{1a}$  receptor, and pharmaceutically acceptable salts thereof.

According to a second aspect, the present invention relates to pharmaceutical compositions comprising these novel compounds, which compositions are useful for the treatment of, *inter alia*, male erectile dysfunction and primary  
30 dysmenorrhoea.

According to a third aspect, the present invention relates to the use of these novel compounds for the manufacture of a pharmaceutical composition for the treatment of erectile dysfunction.


According to a fourth aspect, the present invention relates to the use of  
35 the compounds in the above mentioned series of potent and specific OT recep-

tor agonists and/or VP receptor antagonists for the manufacture of a pharmaceutical composition for treatment of dysmenorrhoea.

According to further aspects, the present invention relates to the use of the above mentioned compounds and compositions in therapy and to therapeutic methods wherein the above mentioned compounds and compositions are used.

### DETAILED DESCRIPTION OF THE INVENTION

Thus, according to the first aspect, the present invention relates to novel specific OT agonists and/or novel specific V<sub>1a</sub> antagonists. These compounds are benzyl carbamates and ureas having the general formula 1:




In this general formula the substituents R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> are, independently of each other, selected from hydrogen (H), alkyl groups, alkoxy (O-alkyl) groups, and the halogens fluorine (F), chlorine (Cl) and bromine (Br). Preferably, at least one of R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> is H and at least one is not H. More preferably, two of R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> are H, and the other is an alkyl group, an O-alkyl group or a halogen.

The linking group X<sup>1</sup> is selected from oxygen (O) and unsubstituted nitrogen (NH). Preferably, X<sup>1</sup> is NH.


The integer a may be 1 or 2, and the integer b may be 1, 2 or 3. Preferably a is 1 and b is 2 such that this ring is a piperazine.

The substituent R<sup>4</sup> is selected from H, alkyl groups, alkenyl groups, alkynyl groups, optionally substituted phenyl, optionally substituted thienyl, optionally substituted furyl, optionally substituted pyridyl, pyrrolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl and isothiazolyl groups, a group - (CH<sub>2</sub>)<sub>e</sub>R<sup>8</sup>, wherein e is 1, 2, 3 or 4, -CH<sub>2</sub>-CH=CH-CH<sub>2</sub>-R<sup>8</sup>, -CH<sub>2</sub>-C≡C-CH<sub>2</sub>-R<sup>8</sup>, -(CH<sub>2</sub>)<sub>g</sub>-CH(OH)-(CH<sub>2</sub>)<sub>h</sub>-R<sup>8</sup>, wherein g and h are, independently of each other, 1 or 2, -(CH<sub>2</sub>)<sub>i</sub>-O-(CH<sub>2</sub>)<sub>j</sub>-R<sup>8</sup> wherein i and j are, independently of each other, 1 or 2, and




wherein  $\text{R}^8$  is selected from H, F,  $\text{CF}_3$ , alkyl groups, alkenyl groups, alkynyl groups, acyl groups, O-alkyl groups, S-alkyl groups, O-acyl groups, hydroxy-alkyl groups, amino groups such as  $\text{NH}_2$ , NH-alkyl,  $\text{N}(\text{alkyl})_2$ , 1-pyrrolidinyl, 1-piperidinyl and 4-morpholinyl, NH-acyl,  $\text{N}(\text{alkyl})$ -acyl,  $\text{CO}_2\text{H}$ ,  $\text{CO}_2$ -alkyl,  $\text{CONH}_2$ ,  $\text{CONH}$ -alkyl,  $\text{CON}(\text{alkyl})_2$ , CN and optionally substituted phenyl, optionally substituted pyridyl, optionally substituted thienyl, optionally substituted furyl, optionally substituted pyrrolyl, optionally substituted pyrazolyl, optionally substituted imidazolyl, optionally substituted oxazolyl, optionally substituted isoxazolyl, optionally substituted thiazolyl and optionally substituted isothiazolyl groups. Suitable optional substituents for the phenyl, pyridyl, thienyl, furyl, pyrrolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl and isothiazolyl groups in  $\text{R}^4$  and  $\text{R}^8$  include F, Cl, Br,  $\text{CF}_3$ , alkyl groups, OH, O-alkyl groups, hydroxyalkyl groups, amino groups such as  $\text{NH}_2$ , NH-alkyl and  $\text{N}(\text{alkyl})_2$ , NH-acyl,  $\text{N}(\text{alkyl})$ -acyl,  $\text{CO}_2\text{H}$ ,  $\text{CO}_2$ -alkyl,  $\text{CONH}_2$ ,  $\text{CONH}$ -alkyl,  $\text{CON}(\text{alkyl})_2$ , oxadiazolyl, thiadiazolyl, CN and  $\text{NO}_2$ . The phenyl, pyridyl, thienyl, furyl, pyrrolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl or isothiazolyl group may have up to three such substituents which may be the same or different.


The group  $\text{G}^1$  is a disubstituted nitrogen such that the  $\text{C}(=\text{O})-\text{G}^1$  bond is an amide bond.  $\text{G}^1$  is selected from an acyclic group according to general formula 2, a fused bicyclic group according to general formulae 3, 4 and 5, and a fused tricyclic group according to general formulae 6 and 7.




2



3




4




5

8



6



7

In general formula 2,  $R^5$  and  $R^6$  are, independently of each other, selected from alkyl, Ar and  $-(CH_2)_f-Ar$ , wherein  $f$  is 1, 2 or 3 and Ar is selected from optionally substituted thienyl and optionally substituted phenyl. Suitable substituents for the phenyl group are alkyl groups, OH, alkoxy groups, halogens,  $NH_2$ ,  $NH$ -alkyl and  $N(alkyl)_2$ . The phenyl group may be substituted with up to three such substituents which may be the same or different.

In general formula 3,  $A^1$  is selected from  $CH_2$ ,  $CH(OH)$ ,  $NH$ ,  $N$ -alkyl, O and S.  $A^2$  is selected from  $CH_2$ ,  $CH(OH)$ ,  $C(=O)$  and  $NH$ , and  $c$  is 1 or 2, preferably 2. It is preferred that when  $A^2$  is  $NH$  then  $A^1$  is  $CH_2$ . It is also preferred that when  $A^2$  is  $C(=O)$  then  $A^1$  is  $NH$  or  $N$ -alkyl.

In general formulae 3, 6 and 7,  $A^3$  is selected from S,  $NH$ ,  $N$ -alkyl,  $-CH=CH-$  and  $-CH=N-$  and  $A^4$ , and  $A^5$  are each, independently of each other, selected from CH and N. In a preferred embodiment,  $A^3$  is S and  $A^4$  and  $A^5$  are both CH, so as to form a thiophene ring. In another preferred embodiment,  $A^3$  is  $-CH=CH-$  and  $A^4$  and  $A^5$  are both CH, so as to form a benzene ring. In another preferred embodiment,  $A^3$  is  $-CH=N-$  and  $A^4$  and  $A^5$  are both CH, so as to form a pyridine ring. In another preferred embodiment,  $A^3$  is  $-CH=CH-$ ,  $A^4$  is CH and  $A^5$  is N, again so as to form a pyridine ring.

In general formulae 4 and 6,  $A^6$  is selected from  $CH_2$ ,  $NH$ ,  $N$ -alkyl and O,  $A^7$  and  $A^{11}$  are, independently of each other, selected from C and N,  $A^8$  and  $A^9$  are, independently of each other, selected from CH, N,  $NH$ ,  $N-(CH_2)_d-R^7$  and S, and  $A^{10}$  is selected from  $-CH=CH-$ , CH, N,  $NH$ ,  $N-(CH_2)_d-R^7$  and S, wherein  $d$  is 1, 2 or 3 and  $R^7$  is selected from H, F,  $CF_3$ , alkyl groups, OH, O-alkyl groups, S-alkyl groups, O-acyl groups, amino groups such as  $NH_2$ ,  $NH$ -alkyl and  $N(alkyl)_2$ ,  $NH$ -acyl,  $N(alkyl)$ -acyl,  $CO_2H$ ,  $CO_2$ -alkyl,  $CONH_2$ ,  $CONH$ -alkyl,  $CON(alkyl)_2$ , CN and optionally substituted phenyl groups. Suitable optional substituents for the phenyl groups in  $R^7$  include F, Cl, Br,  $CF_3$ , alkyl groups, O-alkyl groups, amino groups such as  $NH_2$ ,  $NH$ -alkyl and  $N(alkyl)_2$ ,  $NH$ -acyl,  $N(alkyl)$ -acyl,  $CO_2H$ ,  $CO_2$ -alkyl,  $CONH_2$ ,  $CONH$ -alkyl,

CON(alkyl)<sub>2</sub>, CN and NO<sub>2</sub>. The phenyl group may have up to three such substituents which may be the same or different.

The ring constituted by A<sup>7</sup>, A<sup>8</sup>, A<sup>9</sup>, A<sup>10</sup> and A<sup>11</sup> is aromatic, and accordingly the groups must satisfy certain requirements. When A<sup>10</sup> is -CH=CH- the ring is a six-membered ring. As such, it can only comprise atoms of the type -C(R)= and -N=. Hence A<sup>7</sup> and A<sup>11</sup> must both be C and A<sup>8</sup> and A<sup>9</sup> must be either CH or N. When A<sup>10</sup> is not -CH=CH- then the ring is a five-membered ring. In this case one, and only one, of the atoms in the ring must be S or a trigonal nitrogen. In this context, a "trigonal nitrogen" is a nitrogen atom linked covalently to three different atoms. Two of these atoms are the immediate neighbours to the nitrogen atom in the five-membered ring. The third is a hydrogen, carbon or other atom linked to the five-membered ring. Thus it follows that, when A<sup>10</sup> is not -CH=CH- then one (and only one) of A<sup>7</sup>, A<sup>8</sup>, A<sup>9</sup>, A<sup>10</sup> and A<sup>11</sup> must be S or a trigonal nitrogen. Hence the selection of A<sup>7</sup>, A<sup>8</sup>, A<sup>9</sup>, A<sup>10</sup> and A<sup>11</sup> is subject to the following restrictions.

- 1) If A<sup>10</sup> is not -CH=CH- then one of A<sup>8</sup>, A<sup>9</sup> and A<sup>10</sup> is NH, N-(CH<sub>2</sub>)<sub>d</sub>-R<sup>7</sup> or S or one of A<sup>7</sup> and A<sup>11</sup> is N.
- 2) Not more than one of A<sup>8</sup>, A<sup>9</sup> and A<sup>10</sup> may be NH, N-(CH<sub>2</sub>)<sub>d</sub>-R<sup>7</sup> or S.
- 3) A<sup>7</sup> and A<sup>11</sup> may not both simultaneously be N.
- 4) Neither A<sup>7</sup> nor A<sup>11</sup> may be N if one of A<sup>8</sup>, A<sup>9</sup> and A<sup>10</sup> is NH, N(CH<sub>2</sub>)<sub>d</sub>-R<sup>7</sup> or S

In a preferred embodiment, A<sup>6</sup> is NH. In another preferred embodiment, A<sup>8</sup> is NH or N-(CH<sub>2</sub>)<sub>d</sub>-R<sup>7</sup>. In an even more preferred embodiment, A<sup>8</sup> is NH or N-(CH<sub>2</sub>)<sub>d</sub>-R<sup>7</sup>, and A<sup>9</sup> is N and A<sup>10</sup> is CH.

In general formulae 5 and 7, A<sup>12</sup> and A<sup>13</sup> are selected from N and C and A<sup>14</sup>, A<sup>15</sup> and A<sup>16</sup> are selected from NH, N-CH<sub>3</sub>, S, N and CH. Again, these atoms constitute an aromatic five-membered ring and so there must be one, and only one, S or trigonal nitrogen. Hence the selection of A<sup>12</sup>, A<sup>13</sup>, A<sup>14</sup>, A<sup>15</sup> and A<sup>16</sup> is subject to the following restrictions.

- 1) One of A<sup>14</sup>, A<sup>15</sup> and A<sup>16</sup> is NH, N-CH<sub>3</sub> or S or one of A<sup>12</sup> and A<sup>13</sup> is N.
- 2) Not more than one of A<sup>14</sup>, A<sup>15</sup> and A<sup>16</sup> is NH, N-CH<sub>3</sub> or S.
- 3) A<sup>12</sup> and A<sup>13</sup> may not both simultaneously be N.
- 4) If one of A<sup>14</sup>, A<sup>15</sup> and A<sup>16</sup> is NH, N-CH<sub>3</sub> or S then A<sup>12</sup> and A<sup>13</sup> are both C

As used herein, the term "alkyl" or "alkyl group" is intended to designate lower alkyl groups, i.e. saturated hydrocarbon groups of between one and six carbon atoms, including linear, branched and cyclic alkyl groups. Examples of

"alkyl" include, but are not limited to: C<sub>1</sub> - methyl, C<sub>2</sub> - ethyl, C<sub>3</sub> - propyl, isopropyl, cyclopropyl, C<sub>4</sub> - n-butyl, sec-butyl, isobutyl, tert-butyl, cyclobutyl, cyclopropylmethyl, methylcyclopropyl, C<sub>5</sub> - n-pentyl, neopentyl, cyclopropylethyl, dimethylcyclopropyl, and C<sub>6</sub> - n-hexyl, cyclohexyl, bicyclo[3.1.0]hexyl.

The term "alkenyl" or "alkenyl group" denotes a lower alkenyl group, i.e. a mono-unsaturated hydrocarbon group of between two and six carbon atoms, including linear, branched and cyclic alkenyl groups. Examples of "alkenyl" include, but are not limited to: C<sub>2</sub> - vinyl, C<sub>3</sub> - allyl, 1-methylvinyl, 1-propenyl, C<sub>4</sub> - but-3-enyl, but-2-enyl, methallyl.

The term "alkynyl" or "alkynyl group" denotes a lower alkynyl group, i.e. an unsaturated hydrocarbon group of between two and six carbon atoms which includes a carbon-carbon triple bond, including linear, branched and cyclic alkynyl groups. Examples of "alkynyl" include, but are not limited to: C<sub>2</sub> - ethynyl, C<sub>3</sub> - propargyl, 1-propynyl.

The term "hydroxyalkyl" denotes an alkyl group as defined above in which one or more of the hydrogen atoms are replaced by hydroxyl groups (OH). In general, not more than one hydroxyl group will be attached to any particular carbon atom within the hydroxyalkyl group. Examples of hydroxyalkyl groups include, but are not limited to: hydroxymethyl (HOCH<sub>2</sub>), 1-hydroxyethyl (CH<sub>3</sub>CH(OH)), 2-hydroxyethyl (HOCH<sub>2</sub>CH<sub>2</sub>), 1,2-dihydroxyethyl (HOCH<sub>2</sub>CH(OH)), 4-hydroxy-2-pentyl (CH<sub>3</sub>CH(OH)CH<sub>2</sub>CH(CH<sub>3</sub>)), and 4-hydroxycyclohexyl.

The term "acyl" denotes a group R-C(=O), where R is H, a saturated or unsaturated hydrocarbon moiety of up to seven carbon atoms or a pyridyl or thienyl group. Examples of acyl groups include, but are not limited to: formyl, acetyl, pivaloyl, benzoyl and nicotinoyl.

The compounds according to the present invention generally contain a basic nitrogen atom and so are capable of forming addition salts with protic acids such as hydrochloric acid, sulphuric acid, phosphoric acid, acetic acid, trifluoroacetic acid, benzoic acid, maleic acid, citric acid, fumaric acid, methanesulphonic acid and the like. The compounds of the present invention may also contain an acidic group, such as a carboxylic acid group at R<sup>7</sup> or R<sup>8</sup>. These compounds may exist as inner salts (zwitterions) or as salts such as sodium, potassium, magnesium, calcium or tetra-alkylammonium salts. To the




extent that such salts are pharmaceutically acceptable, they are included within the scope of the present invention.

The compounds according to the present invention may have one or more stereogenic centres ("asymmetric carbon atoms") and so may exhibit optical isomerism. The scope of the present invention includes all epimers, enantiomers and diastereomers of compounds according to general formula 1, including single isomers, mixtures and racemates.

Particularly preferred embodiments within the present invention are those compounds that combine two or more of the preferred features described above.


One such particularly preferred embodiment is a urea according to general formula 8.



8

In general formula 8,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$  and  $G^1$  are as previously defined.

Another particularly preferred embodiment is a compound according to general formula 9, which corresponds to a compound according to general formula 1 in which  $G^1$  is a group according to general formula 6 wherein  $A^4$ ,  $A^5$  and  $A^{10}$  are all CH,  $A^6$  is NH,  $A^7$  and  $A^{11}$  are both C,  $A^8$  is  $N(CH_2)_dR^7$  and  $A^9$  is N.




9

12

In general formula 9,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^7$ ,  $A^3$ ,  $X^1$ ,  $a$ ,  $b$  and  $d$  are as previously defined.

A most preferred embodiment is a compound according to general formula 10.



10

5 In general formula 10,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^7$ ,  $A^3$  and  $d$  are as previously defined.

Individual preferred compounds within the invention include:

4-(3,3-Dimethyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

10 4-(2-Cyclopropyl-ethyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

4-Cyclopropylmethyl-piperazine-1-carboxylic acid 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

15 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 3-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

4-(2-Hydroxymethyl-cyclopropylmethyl)-piperazine-1-carboxylic acid-2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

20 4-(3-Methyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

4-Cyclopentylmethyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

5 4-Cyclohexylmethyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

4-Cyclopropylmethyl-piperazine-1-carboxylic acid 3-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

10 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

4-Cyclobutylmethyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

15 4-(2-Cyclopropyl-ethyl)-piperazine-1-carboxylic acid 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

4-Pentyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

20 4-Hexyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

(R)-4-(2-Methyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

25 4-(2-Ethyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

4-(2-Methyl-but-2-enyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

30 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

35 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 2-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

4-Cyclopropylmethyl-piperazine-1-carboxylic acid 2-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

4-Cyclobutylmethyl-piperazine-1-carboxylic acid 4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

4-Cyclopropylmethyl-piperazine-1-carboxylic acid 2-ethyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

4-Cyclobutylmethyl-piperazine-1-carboxylic acid 2-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

4-Cyclopropylmethyl-piperazine-1-carboxylic acid 2-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide; and

4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-methoxy-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide.

Other individual preferred compounds, especially for treatment of primary dysmenorrhoea, and also for treatment of pre-term labour, Raynaud's disease, brain oedema, motion sickness, small cell lung cancer, depression, anxiety, hyponatremia, liver cirrhosis or congestive heart failure, within the invention include:


4-Cyclopropylmethyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide; and

4-(3-Methylsulfanyl-propyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide.

These individually preferred compounds have several advantages over existing entities. They are selective for the  $V_{1a}$  receptor over other AVP receptors. Thus they are potentially safer and more effective than existing non-selective entities. Existing  $V_{1a}$  selective entities have been suspended during development due to issues such as safety. Furthermore, they are non-peptidic small molecules. It is well known that such compounds have significantly more potential to be orally active than peptides. As such they offer more convenience and better patient compliance than peptidic entities.

The compounds of the present invention can be prepared by standard chemical manipulations. In general, compounds according to general formula 1 can be considered to consist of three component parts:

- Component C<sup>1</sup> corresponding to G<sup>1</sup>
- 5    • Component C<sup>2</sup> corresponding to the substituted benzoyl unit
- Component C<sup>3</sup> corresponding to the saturated heterocycle




Intermediates corresponding to these components are prepared and then assembled to give the final product. These three components are:


- (i) for C<sup>1</sup>, a secondary amine



- (ii) for C<sup>2</sup>, a substituted benzoic acid



- (iii) for C<sup>3</sup>, a monosubstituted saturated heterocycle




It will be recognised that the substituted benzoic acid that serves for C<sup>2</sup> has two functional groups, one of which will need temporary protection during the assembly of the final compound. The principles of functional group protection are well known in the art and are described in, for example, J.F.W. McOmie, "Protective Groups in Organic Chemistry", Plenum Press, 1973; T.W. Greene and P.G.M. Wuts, "Protective Groups in Organic Synthesis", 2<sup>nd</sup> edition, John Wiley, 1991; and P.J. Kocienski, "Protecting groups", Georg Thieme Verlag, 1994. The carboxylic acid group will usually be protected as

an ester, such as the methyl, benzyl or *tert*-butyl ester. The primary amine of the benzoic acid (when  $X^1 = \text{NH}$ ) will usually be protected as a carbamate derivative such as the *tert*-butyl carbamate (BOC derivative), the benzyl carbamate (CBZ or more simply Z derivative) or the 9-fluorenylmethyl carbamate (Fmoc derivative). When  $X^1 = \text{O}$  the resulting alcohol function will usually be protected as an ester such as an acetate, or an ether such as a methoxymethyl, tetrahydropyranyl or trialkylsilyl ether. Other functional groups may require protection. For example, the group  $G^1$  may include one or more primary or secondary amino groups which may need protection. In the following general description of the synthetic methodology it will be assumed that such protection is used when necessary.

*(i) Preparation of secondary amine for  $C^1$*


Acyclic secondary amines corresponding to  $\text{HNR}^5\text{R}^6$  are well known. Many are items of commerce. Those that are not may be prepared according to published methods or by simple modification of such methods. Some particularly useful methods are listed below.

a) Alkylation




(This method is only applicable in cases where further alkylation can be avoided.)

b) Reductive amination



(where  $\text{R}^a\text{CHR}^b$  corresponds to  $\text{R}^6$ )


c) Amide reduction



17

(where  $R^aCH_2$  corresponds to  $R^6$ )

The starting amide can itself be prepared using well known methods.




- 5 Secondary amines corresponding to  $C^1$  where  $G^{11}$  is a group according to general formulae 3 – 7 are generally not commercially available. They can be prepared according to published methods, or by obvious modifications of such methods. Particularly useful methods are described in: Aranapakam *et al.*, Bio-  
 10 org. Med. Chem. Lett. 1993, 1733; Artico *et al.*, Farmaco. Ed. Sci. 24, 1969, 276; Artico *et al.*, Farmaco. Ed. Sci. 32, 1977, 339; Chakrabarti *et al.*, J. Med. Chem. 23, 1980, 878; Chakrabarti *et al.*, J. Med. Chem. 23, 1980, 884; Chakrabarti *et al.*, J. Med. Chem. 32, 1989, 2573; Chimirri *et al.*, Heterocycles 36, 1993, 601; Grunewald *et al.*, J. Med. Chem. 39, 1996, 3539; Klunder *et al.*, J. Med. Chem. 35, 1992, 1887; Liegeois *et al.*, J. Med. Chem. 37, 1994, 519;  
 15 Olagbemiro *et al.*, J. Het. Chem. 19, 1982, 1501; Wright *et al.*, J. Med. Chem. 23, 1980, 462; Yamamoto *et al.*, Tet. Lett. 24, 1983, 4711; and International patent application, publication number WO99/06403.

(ii) Preparation of substituted benzoic acid for  $C^2$

- 20 Substituted benzoic acids corresponding to  $C^2$  are not generally items of commerce, but they can be prepared using published methods or obvious variations of such methods. The main challenge is generally the elaboration of the  $-CH_2X^1H$  functionality at the 4-position. Some useful transformations are listed below.


18

## a) Bromination/Substitution



5


## b) Sandmeyer

(iii) Preparation of heterocycle derivative for C<sup>3</sup>

10

Certain heterocycles corresponding to C<sup>3</sup>, particularly *N*-aryl piperazines, are items of commerce. Other heterocycles can be prepared according to the methods described in the literature. Useful transformations include the following.

## a) Alkylation or reductive alkylation




15

(where PG is a protecting group and R<sup>A</sup>CH<sub>2</sub> is R<sup>4</sup>)




19

**b) Acylation/reduction**




### c) Reduction



5

With the three components, suitably protected if necessary, in hand, the assembly of the final compound requires the formation of two bonds: between C<sup>1</sup> and C<sup>2</sup>, and between C<sup>2</sup> and C<sup>3</sup>. These bond-forming steps may be taken in either order. Thus, the following sequences can be proposed:



15 (i) Formation of  $C^1-C^2$  bond

The bond between C<sup>1</sup> and C<sup>2</sup> is a simple amide bond. The chemistry for making such bonds from a carboxylic acid and a secondary amine is well known in the art of organic synthesis, and particularly in the field of peptide synthesis. The carboxylic acid may be converted into a more reactive species such as an acid chloride (using, for example oxalyl chloride or thionyl chloride) or a mixed anhydride (using isobutyl chloroformate). This reactive species is then added to the secondary amine in a suitable solvent, generally an aprotic solvent such as dichloromethane or dimethylformamide, in the presence of a base such as triethylamine or 4-dimethylaminopyridine, and the reaction is allowed to proceed at a temperature between -20°C and the boiling point of the solvent. The choice of temperature and the time allowed for the reaction will depend on the reactivity of the two components.

Alternatively, the carboxylic acid and the secondary amine may be mixed in a suitable solvent as above, optionally in the presence of a base, and a  
30 condensing agent added. Suitable condensing agents include carbodiimides, such as dicyclohexylcarbodiimide (DCC) and N-ethyl-N'-

dimethylaminopropylcarbodiimide (EDC, also WSCD for water-soluble carbodiimide), phosphorus reagents such as (benzotriazol-1-yloxy)-tris(dimethylamino)phosphonium hexafluorophosphate (BOP), (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyBOP<sup>®</sup>) and bromotripyrrolidinophosphonium hexafluorophosphate (PyBroP<sup>®</sup>), and ureas such as O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU).

*(ii) Formation of C<sup>2</sup>-C<sup>3</sup> bond*

10       The bond between C<sup>2</sup> and C<sup>3</sup> is a carbamate (when X<sup>1</sup> = O) or a urea (when X<sup>1</sup> = NH). The first step in the formation of this bond is generally to react the heterocycle derivative with phosgene or a phosgene equivalent such as trichloromethyl chloroformate, bis(trichloromethyl)carbonate or carbonyldiimidazole. Again, an aprotic solvent and a tertiary amine base will generally  
15       be used. The intermediate formed in this step is usually not isolated. The alcohol (X<sup>1</sup> = O) or amine (X<sup>1</sup> = NH) is added and the reaction is allowed to continue, directly forming the carbamate or urea. As an alternative, when X<sup>1</sup> = NH the reactive intermediate may be formed by the reaction of C<sup>2</sup> with the phosgene equivalent and the amine added in the second part of the synthesis.

20       The compounds according to the present invention are useful in human and animal therapy.

As stated above, the second, third and fourth aspect of the present invention relates to pharmaceutical compositions comprising the above described compounds and to the manufacture of pharmaceutical compositions using the  
25       above described compounds. In such pharmaceutical compositions, the above described compounds constitutes a pharmaceutically active ingredient. It may be the sole active ingredient, or it may be combined by at least one other active ingredient or agent. Preferably the pharmaceutical composition includes no additional active agents. Normally, the pharmaceutical compositions according  
30       to the invention, used according to the invention or produced according to the invention also comprise other substances, such as an inert vehicle, or pharmaceutical acceptable adjuvants, carriers, preservatives etc., which are well known to persons skilled in the art.

35       The pharmaceutical composition according to the present invention may be presented in any form that is known in the art. For example, the formulation may be presented as a tablet, capsule, powder, suppository, cream, solution or

suspension, or in a more complex form such as an adhesive patch. The formulation will generally include one or more excipients, such as diluents, bulking agents, binding agents, dispersants, solvents, preservatives, flavouring agents and the like. Where the formulation is presented as a tablet or capsule the excipients may optionally include one or more agents to control the release of the active species, such as a coating of a polymer that is insoluble at low pH but soluble at neutral or high pH. Such a coating (known as an "enteric coating") prevents the release of the active agent in the stomach but allows its release in the intestines.

The compounds according to the present invention are useful for treatment of several diseases, disorders or conditions. The term "treatment" used herein relates to both treatment in order to cure or alleviate a disease, disorder or a condition, and to treatment in order to prevent the development of a disease, disorder or a condition. The treatment may either be performed in an acute or in a chronic way. The human or animal to be treated, i.e. the patient, may be any human or non-human mammal in need of treatment according to the invention.

The novel compounds of the present invention are potent and selective OT agonists and/or  $V_{1a}$  receptor antagonists and so they and pharmaceutical compositions comprising them are useful in the treatment of treatment of conditions for which inadequate oxytocin-like activity is implicated in the pathophysiology and in the treatment of conditions in which inappropriate vasopressin-like activity is implicated in the pathophysiology. Conditions for which inadequate oxytocin-like activity is implicated in the pathophysiology include, but are not limited to: sexual disorders such as male erectile dysfunction, ejaculatory disorders and female sexual dysfunction, cancer of the prostate, breast, ovary and bones, osteoporosis, benign prostatic hyperplasia, postpartum bleeding, and depression. Conditions in which inappropriate vasopressin-like activity is implicated in the pathophysiology include disorders affecting blood platelets, blood vessels, hepatocytes, brain and uterus-cervix. The novel  $V_{1a}$  receptor antagonist and/or pharmaceutical compositions comprising them are suitable for treatment of primary dysmenorrhoea, pre-term labour, hypertension, Raynaud's disease, brain oedema, motion sickness, small cell lung cancer, depression, anxiety, hyponatremia, liver cirrhosis and congestive heart failure.. In a preferred embodiment, the compounds or pharmaceutical

compositions according to the invention are used for treatment of primary dysmenorrhoea.

Further aspects of the invention relates to methods for treatment of the above mentioned diseases, disorders or conditions, and preferably a method for treatment of primary dysmenorrhoea. According to the method according to the invention a therapeutically effective amount of the compound, or of the pharmaceutical composition, described above is administered to a patient in need of this treatment.

The term "therapeutically effective amount" relates to an amount that will lead to the desired therapeutical effect. The therapeutically effective amount will be determined by the attending physician taking into consideration all appropriate factors. Generally a single dose will comprise between 0.1 mg and 1000 mg, preferably between 1 mg and 250 mg, of the active compound according to the invention. The dose may be given on a single occasion or repeatedly. When given repeatedly, it may be given at regular intervals, such as once, twice or three times daily, or on demand, according to the condition being treated.

When used as therapeutic agents, the compositions of the present invention may be administered by any appropriate route that is known in the art. For example, they may be administered by the oral, buccal, sublingual, rectal, intravaginal, nasal, pulmonary or transdermal routes. Alternatively, they may be given by injection, including intravenous, subcutaneous and intramuscular injection.

For long-term treatment an alternative to repeated dosing may be the administration of a depot dose. For this method of administration the active agent is generally introduced into a matrix of biodegradable polymer, such as a copolymer of lactic and glycolic acids, and the formulation is given either subcutaneous (s.c.) or intramuscularly (i.m.) so as to form a deposit from which the active agent is released as the polymer degrades.

In order to decide whether or not a compound having general formula I is a selective  $V_{1a}$  antagonist the compound may be assayed to determine its ability to inhibit the cellular consequences of AVP stimulation on intact cells. In the assay, the compounds cause significant inhibition of cellular activation at concentrations of 30  $\mu$ M or less. Preferred compounds cause significant inhibition at concentrations of 300 nM or less.

The present invention is further illustrated in the following examples, which are intended to demonstrate the application of the invention but not to limit the scope thereof.


## 5 EXAMPLES

The following abbreviations are used:

Bu	butyl – alkyl residues may be further denoted as n (normal, i.e. unbranched), i (iso) and t (tertiary)
DIEA	<i>N,N</i> -diisopropylethylamine
DMF	dimethylformamide
Et	ethyl
EtOAc	ethyl acetate
HOBt	1-hydroxybenzotriazole
HPLC	high pressure liquid chromatography
h	hour(s)
Me	methyl
MS	mass spectrum
NMR	nuclear magnetic resonance spectrum – NMR spectra were recorded in CDCl <sub>3</sub> unless otherwise indicated
OVA	ornithine vasotocin analogue
pet. ether	petroleum ether boiling in the range 60-80°C
Ph	phenyl
Pn	pentyl
Pr	propyl
THF	tetrahydrofuran
WSCD	water-soluble carbodiimide ( <i>N</i> -ethyl- <i>N'</i> -(3-dimethylaminopropyl)-carbodiimide hydrochloride)

- 10 Examples 1 – 13 describe the synthesis of intermediates. Compounds according to the present invention are described in Examples 14 – 149. Example 150 describes how compounds can be assayed based on their ability to inhibit the cellular consequences of AVP stimulation on intact cells. Example 151 describes tablets for oral administration comprising a compound according to the invention.

## Example 1

1-Benzyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine

## 5 1A: Ethyl 5-amino-1-benzylpyrazole-4-carboxylate

Benzylhydrazine dihydrochloride (4.29 g, 22 mmol) was added to a solution of ethyl (ethoxymethylene)cyanoacetate (3.38 g, 20 mmol) and triethylamine (6.15 ml, 44 mmol, 2 eq) in ethanol (40 ml) and the mixture was heated at reflux for 18h. The solvent was removed *in vacuo* and the residue was purified by flash chromatography on silica gel (eluant 60% pet. ether/40% ethyl

acetate) to yield a pale yellow solid identified as ethyl 5-amino-1-benzylpyrazole-4-carboxylate (4.3 g, 88%).

**1B: Ethyl 1-benzyl-5-(2'-nitrophenylamino)pyrazole-4-carboxylate**

- 5 Sodium hydride (60% dispersion in oil, 520 mg, 13 mmol) was added portionwise to a suspension of ethyl 5-amino-1-benzylpyrazole-4-carboxylate (2.2 g, 9 mmol) in anhydrous THF (30ml) at 0°C. The mixture was allowed to warm to room temperature and stirred for 2h then 1-fluoro-2-nitrobenzene (1.26 g, 9 mmol) was added and the resultant deep purple suspension was
- 10 stirred at room temperature for 18 h. 1 M KHSO<sub>4</sub> was added to quench the reaction and the solvent was removed *in vacuo*. The residue was dissolved in ethyl acetate and the solution was washed with 0.3 M KHSO<sub>4</sub>, sat. NaHCO<sub>3</sub> and brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant 75% pet. ether/25% ethyl
- 15 acetate) to yield ethyl 1-benzyl-5-(2'-nitrophenylamino)pyrazole-4-carboxylate (2.5 g, 76%).

MS [M+H]<sup>+</sup> 366.8

**20 1C: Ethyl 5-(2'-aminophenylamino)-1-benzylpyrazole-4-carboxylate**

- Ethyl 1-benzyl-5-(2'-nitrophenylamino)pyrazole-4-carboxylate (2.5 g, 6.8 mmol) was dissolved in ethyl acetate/ethanol (1:1, 100 ml) and hydrogenated over 10% Pd/C catalyst for 70 minutes. The mixture was filtered through Celite<sup>®</sup> filter agent and the filtrate was concentrated *in vacuo* to give a white
- 25 solid identified as ethyl 5-(2'-aminophenylamino)-1-benzylpyrazole-4-carboxylate (1.5 g, 86%).

MS [M+H]<sup>+</sup> 337.2

**30 1D: 1-Benzyl-4,10-dihydropyrazolo[5,4-b][1,5]benzodiazepin-4(5H)-one**

- A solution of ethyl 5-(2'-aminophenylamino)-1-benzylpyrazole-4-carboxylate (1.75 g, 5.2 mmol) in acetic acid/ 2-propanol (1:9, 40 ml) was heated at reflux for 3 days. The solvent was removed *in vacuo* and the residue was azeotroped with toluene to give an off-white solid that was purified by
- 35 flash chromatography on silica gel (eluant 35% pet. ether/65% ethyl acetate) to

yield a white solid identified as 1-benzyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepin-4(5*H*)-one (780 mg, 52%).

MS [M+H]<sup>+</sup> 291.1

5

**1E: 1-Benzyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine**

LiAlH<sub>4</sub> (365 mg, 10 mmol) was added portionwise to a suspension of 1-benzyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepin-4(5*H*)-one (780 mg, 2.7 mmol) in anhydrous THF (15 ml) at 0°C over 10 min. The resulting suspension was heated at reflux for 18 h, then allowed to cool to room temperature. A further portion of LiAlH<sub>4</sub> (90 mg, 2.5 mmol) was added and the mixture was heated at refluxed for 3 h. The mixture was cooled to 0°C, 35% ammonia solution (1 ml) was added dropwise over 10 min and the mixture was stirred at room temperature for 1h. The resulting suspension was filtered through Celite<sup>®</sup> filter agent and the filtrate was concentrated *in vacuo* to give a white solid identified as 1-benzyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine (450 mg, 60%).


10

15

MS [M+H]<sup>+</sup> 276.9

20



**Example 2****1-Methyl-4,10-dihydropyrazolo[4,5-c]pyrido[2,3-b][1,4]diazepine****2A: Ethyl 1-methyl-2-(3'-nitro-2'-pyridylamino)pyrazole-4-carboxylate**

5 Sodium hydride (60% dispersion in oil, 600 mg, 15mmol) was added portionwise to a suspension of ethyl 5-amino-1-methylpyrazole-4-carboxylate (1.69 g, 10 mmol) in anhydrous THF (15ml) at 0°C. The mixture was stirred for 2h at room temperature then 2-chloro-3-nitropyridine (1.58 g, 10 mmol) was added and the resulting deep red suspension was stirred at room temperature for 18 h. 1 M KHSO<sub>4</sub> was added to quench the reaction and the solvent was removed *in vacuo*. The residue was dissolved in ethyl acetate and the solution was washed with 0.3 M KHSO<sub>4</sub>, sat. NaHCO<sub>3</sub> and brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant 30% pet. ether/70% ethyl acetate) to give ethyl 1-methyl-2-(3'-nitro-2'-pyridylamino)pyrazole-4-carboxylate (1.95 g, 67%).

15

MS [M+H]<sup>+</sup> 292.0

**2B: Ethyl 2-(3'-amino-2'-pyridylamino)-1-methylpyrazole-4-carboxylate**

A solution of ethyl 1-methyl-2-(3'-nitro-2'-pyridylamino)pyrazole-4-carboxylate (1.95 g, 6.7 mmol) in ethanol (100ml) was hydrogenated over 10% Pd/C catalyst for 3 h. The reaction mixture was filtered through Celite® filter agent and the filtrate was concentrated *in vacuo* to give a white solid identified as ethyl 2-(3'-amino-2'-pyridylamino)-1-methylpyrazole-4-carboxylate (1.5 g, 86%).

**2C: 1-Methyl-4,10-dihydropyrazolo[4,5-*c*]pyrido[2,3-*b*][1,4]diazepin-4(5*H*)-one**

A solution of ethyl 2-(3'-amino-2'-pyridylamino)-1-methylpyrazole-4-carboxylate (1.5 g, 5.75 mmol) in acetic acid/2-propanol (1:9, 50 ml) was heated at reflux for 3 days. The solvent was removed *in vacuo* and the residue was azeotroped with toluene. The residue was purified by recrystallization from ethanol and then flash chromatography on silica gel (eluant 95% chloroform/4% methanol/1% acetic acid) to give a white solid identified as 1-methyl-4,10-dihydropyrazolo[4,5-*c*]pyrido[2,3-*b*][1,4]diazepin-4(5*H*)-one (560 mg, 45%).


20

**2D: 1-Methyl-4,10-dihydropyrazolo[4,5-*c*]pyrido[2,3-*b*][1,4]diazepine**

LiAlH<sub>4</sub> (365 mg, 10 mmol) was added portionwise to a suspension of 1-methyl-4,10-dihydropyrazolo[4,5-*c*]pyrido[2,3-*b*][1,4]diazepin-4(5*H*)-one (560 mg, 2.6 mmol) in anhydrous THF (30 ml) at 0°C over 10 minutes. The resulting suspension was heated at reflux for 18 h. The reaction was cooled to 0°C and 35% ammonia solution (1 ml) was added dropwise over 10 minutes, then the mixture was stirred at room temperature for 1 h. The resulting suspension was filtered through Celite® filter agent and the filtrate was concentrated *in vacuo* to give a white solid identified as 1-methyl-4,10-dihydropyrazolo[4,5-*c*]pyrido[2,3-*b*][1,4]diazepine (410 mg, 78%).

30

MS [M+H]<sup>+</sup> 202.1.

**Example 3*****tert*-Butyl 4-aminomethyl-3-chlorobenzoate****5 3A: *tert*-Butyl 3-chloro-4-methylbenzoate**

Thionyl chloride (11 ml, 150 mmol) was added to a suspension of 3-chloro-4-methylbenzoic acid (5.12 g, 30 mmol) in toluene (25 ml) and the mixture was heated at reflux for 2 h. The solvent was removed *in vacuo* and the residue was azeotroped with toluene three times, then dissolved in anhydrous THF (40 ml) and cooled to 0°C. Lithium *tert*-butoxide (2.4 g, 30 mmol) was added and the mixture was stirred at room temperature for 3 days. Water (5 ml) was added and the solvent was removed *in vacuo*. The residue was dissolved in ethyl acetate. The solution was washed with 0.3M KHSO<sub>4</sub>, sat. NaHCO<sub>3</sub> and brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated *in vacuo* to give a pale yellow gum identified as *tert*-butyl 3-chloro-4-methylbenzoate (5.4 g, 79%).

**3B: *tert*-Butyl 4-bromomethyl-3-chlorobenzoate**

*N*-Bromosuccinimide (4.27 g, 24 mmol) and 2,2'-azo-bis(2-methylpropionitrile) (394 mg, 2.4 mmol) were added to a solution of *tert*-butyl 3-chloro-4-methylbenzoate (5.4 g, 23.8 mmol) in carbon tetrachloride (75 ml) and the mixture was heated at reflux for 18 h. The solvent was removed *in vacuo* and the residue was purified by flash chromatography on silica gel (eluant 95% pet.ether/5% ethyl acetate) to give a white solid identified as *tert*-butyl 4-bromomethyl-3-chlorobenzoate (5.7 g, 78%).

25


**3C: *tert*-Butyl 4-aminomethyl-3-chlorobenzoate**

Ethanol (100 ml) was saturated with ammonia, then *tert*-butyl 4-bromomethyl-3-chlorobenzoate (5.7 g, 18.7 mmol) was added and the mixture was stirred at room temperature for 2 h. The solvent was removed *in vacuo* and

the residue was triturated with diethyl ether to give a white solid identified as *tert*-butyl 4-aminomethyl-3-chlorobenzoate (4.1 g, 91%).

#### Example 4

##### 5 4-(*tert*-Butyloxycarbonylaminomethyl)-3-chlorobenzoic acid



##### 4A. Methyl 4-bromomethyl-3-chlorobenzoate

To a solution of methyl 3-chloro-4-methylbenzoate (5.0 g, 27.1 mmol) in carbon tetrachloride (50 ml) were added *N*-bromosuccinimide (5.8 g, 32.0 mmol) and 2,2'-azo-*bis*(2-methylpropionitrile) (0.442 g, 2.70 mmol). The mixture was heated at reflux for 18 h, then allowed to cool to room temperature and concentrated *in vacuo*. The residue was purified by flash chromatography on silica (eluant pet. ether → 5% ethyl acetate/95% pet. ether) to give an oil identified as methyl 4-bromomethyl-3-chlorobenzoate (5.96 g, 84%).


15

##### 4B. 4-(*tert*-Butyloxycarbonylaminomethyl)-3-chlorobenzoic acid


To a saturated solution of ammonia in ethanol (170 ml) was added methyl 4-bromomethyl-3-chlorobenzoate from Example 4A (5.5 g, 20.9 mmol). The mixture was stirred at room temperature for 1 h and then concentrated *in vacuo*. The residue was triturated with diethyl ether and the resultant white crystals were filtered off and washed with more diethyl ether. To a solution of this solid in water (100 ml) were added solutions of di-*tert*-butyl dicarbonate (5.0 g, 23.0 mmol) in dioxan (100 ml) and sodium hydroxide (1.86 g, 46.0 mmol) in water (100 ml). The mixture was stirred at room temperature for 18 h and then concentrated *in vacuo*. The aqueous residue was acidified with citric acid and extracted with chloroform/2-propanol. The organic layer was washed with water, dried over MgSO<sub>4</sub>, and concentrated *in vacuo* to give a white solid identified as 4-(*tert*-butyloxycarbonylaminomethyl)-3-chlorobenzoic acid (2.8 g, 67%).

30

31.


**Example 5****4-(*tert*-Butyloxycarbonylaminomethyl)-3-nitrobenzoic acid**

4-Bromomethyl-3-nitrobenzoic acid (4.75 g, 18.2 mmol) was reacted  
5 following the method of Example 4B to give a yellow solid identified as 4-  
(*tert*-butyloxycarbonylaminomethyl)-3-nitrobenzoic acid (2.6 g, 49%).


**Example 6****4-Cyano-3-methylbenzoic acid**

10

To a solution of 4-bromo-2-methylbenzonitrile (2.0 g, 10.2 mmol) in  
THF (100 ml) at -78°C under a nitrogen atmosphere was added dropwise a 2.5  
M solution of *n*-butyl lithium (4.48 ml, 11.2 mmol). The mixture was stirred at  
-78°C for 1h and then poured onto solid carbon dioxide (5 g) in THF (50 ml).  
15 The mixture was allowed to warm to room temperature. Water was added (200  
ml) and the mixture was extracted with diethyl ether (3 times). The aqueous  
layer was acidified by addition of concentrated HCl and extracted with chloro-  
form (3 times). The combined chloroform extracts were washed with water,  
dried over MgSO<sub>4</sub>, and concentrated *in vacuo* to give a white solid identified as  
20 4-cyano-3-methylbenzoic acid (1.2 g, 73%).

**Example 7****4-Cyano-2-methylbenzoic acid**

4-Bromo-3-methylbenzonitrile (2.0 g, 10.2 mmol) was reacted following the method of Example 6. The product was triturated with hexane to give a yellow solid identified as 4-cyano-2-methylbenzoic acid (0.96 g, 59%).

**Example 8****4-(*tert*-Butyloxycarbonylaminomethyl)-2-fluorobenzoic acid**

10

**8A. 2-Fluoro-4-methylbenzoic acid**

4-Bromo-3-fluorotoluene (8.33 g, 44.07 mmol) was reacted following the method of Example 6 to give a white solid identified as 2-fluoro-4-methylbenzoic acid (4.89 g, 72%).

15

**8B. Methyl 2-fluoro-4-methylbenzoate**


To a solution of 2-fluoro-4-methylbenzoic acid (6.04 g, 39.18 mmol) in toluene (80 ml) was added thionyl chloride (65 ml, 89.11 mmol). The mixture was heated at reflux for 2.5 h, cooled and concentrated *in vacuo*. The residue was dissolved in dichloromethane (50ml) and methanol (50 ml) was added. The mixture was stirred at room temperature for 2.5 h and then concentrated *in vacuo*. The residue was dissolved in dichloromethane (100 ml), washed with saturated sodium bicarbonate solution and brine, dried over  $\text{MgSO}_4$ , and concentrated *in vacuo* to give a tan solid identified as methyl 2-fluoro-4-methylbenzoate (5.07 g, 77%).

**8C. Methyl 4-bromomethyl-2-fluorobenzoate**

Methyl 2-fluoro-4-methylbenzoate (5.07 g, 30.16 mmol) was reacted following the method of Example of 4A. The product was purified by flash chromatography on silica (eluant 20% ethyl acetate/ 80% pet. ether) to give an oil identified as methyl 4-bromomethyl-2-fluorobenzoate (5.9 g, 80%).

**8D. 4-(tert-Butyloxycarbonylaminomethyl)-2-fluorobenzoic acid**

Methyl 4-bromomethyl-2-fluorobenzoate (5.9 g, 24.13mmol) was reacted following the method of Example 4B. The product was recrystallised from dioxan/pet. ether to give white crystals identified as 4-(tert-butyloxycarbonylaminomethyl)-2-fluorobenzoic acid (2.4 6g, 38%).

**Example 9****25 4-Cyano-3,5-dimethylbenzoic acid****9A. 4-Bromo-2,6-dimethylbenzonitrile**

4-Bromo-2,6-dimethylaniline (4.49 g, 22.4mmol) was taken up in water (25ml) and concentrated hydrochloric acid (8.0 ml) was added. The mixture was sonicated to form a fine suspension and then cooled to  $0^\circ\text{C}$ . A solution of sodium nitrite (1.67 g, 24.2mmol) in water (5 ml) was then added dropwise so


as to maintain the temperature of the reaction between 0-5°C. The mixture was stirred at 0-5°C for 30 minutes and then neutralised by addition of solid sodium bicarbonate. The resulting solution was then added portionwise to a solution of copper cyanide (2.42 g, 27.0 mmol) and potassium cyanide (3.65 g, 56.1 mmol) in water (25 ml) at 70°C. The mixture was stirred at 70°C for 30 minutes, allowed to cool and then extracted with toluene (2 times). The combined extracts were washed with water and brine, dried over MgSO<sub>4</sub>, and concentrated *in vacuo*. The residue was purified by flash chromatography on silica (eluant 5% ethyl acetate/ 95% pet. ether) to give an orange solid identified as 4-bromo-2,6-dimethylbenzonitrile (3.2 g, 68%).

#### 9B. 4-Cyano-3,5-dimethylbenzoic acid

4-Bromo-2,6-dimethylbenzonitrile (3.20 g, 15.2 mmol) was reacted following the method of Example 6 to give a tan solid identified as 4-cyano-3,5-dimethylbenzoic acid (1.5 g, 56%).

#### Example 10

##### 2-Chloro-4-cyanobenzoic acid



#### 10A. 4-Amino-2-chloro-benzoic acid methyl ester

Acetyl chloride (2.5 ml) was added drop-wise to a solution of 2-chloro-4-cyano-benzoic acid (2.22 g, 12.94 mmol) in methanol (75 ml) while stirring. The mixture was heated at reflux for 18 h, cooled and concentrated *in vacuo*. The residue was taken up in ethyl acetate, washed with saturated NaHCO<sub>3</sub> and brine and concentrated *in vacuo* to give a beige solid identified as 4-amino-2-chloro-benzoic acid methyl ester (2.32 g, 97%).


#### 10B. 2-Chloro-4-cyano-benzoic acid methyl ester

4-Amino-2-chloro-benzoic acid methyl ester (5.00 g, 26.94 mmol) was reacted following the method of Example 9A to give a pale orange solid identified as 2-chloro-4-cyano-benzoic acid methyl ester (2.62 g, 50%).



**10C. 2-Chloro-4-cyanobenzoic acid**

Lithium hydroxide (1.12 g, 26.69 mmol) was added to a solution of 2-chloro-4-cyano-benzoic acid methyl ester (2.60 g, 13.29 mmol) in dioxan/water (4:1, 100ml). The mixture was stirred at room temperature for 3h and concentrated *in vacuo*. The residue was partitioned between 1N hydrochloric acid and chloroform and the organic layer was washed with brine and concentrated *in vacuo*. The residue was recrystallised from a mixture of dioxan and pet. ether to give a pale orange solid identified as 2-chloro-4-cyanobenzoic acid (2.33 g, 97%).

**Example 11****4-(tert-Butoxycarbonylamino-methyl)-3-fluorobenzoic acid**

**11A. 3-Fluoro-4-methylbenzoic acid methyl ester**

3-Fluoro-4-methylbenzoic acid (5.0 g, 32.4 mmol) was reacted following the method of Example 8B to give a brown oil identified as 3-fluoro-4-methylbenzoic acid methyl ester (4.5 g, 83%).

5

**11B. 4-Bromomethyl-3-fluorobenzoic acid methyl ester**

3-Fluoro-4-methylbenzoic acid methyl ester (4.5 g, 26.6 mol) was reacted following the method of Example 4A to give a yellow oil identified as 4-bromomethyl-3-fluorobenzoic acid methyl ester (2.7 g, 41%).

10

**11C. 4-Azidomethyl-3-fluorobenzoic acid methyl ester**

Sodium azide (609mg) was added to a solution of 4-bromomethyl-3-fluorobenzoic acid methyl ester (2.1 g, 8.5 mmol) in DMF (30ml). The mixture was stirred for 18h, diluted with ethyl acetate, washed with water and brine and concentrated *in vacuo* to give a colourless oil identified as 4-azidomethyl-3-fluorobenzoic acid methyl ester (1.78 g, 100%).

15

**11D. 4-Aminomethyl-3-fluorobenzoic acid methyl ester**

Hydrogen was passed through a degassed solution of 4-azidomethyl-3-fluorobenzoic acid methyl ester (2.11 g, 10 mmol) in methanol containing 10% palladium on carbon for 2 h. The reaction mixture was filtered through Celite<sup>®</sup> and the filtrate was concentrated *in vacuo* to give a colourless oil identified as 4-aminomethyl-3-fluorobenzoic acid methyl ester (1.51 g, 83%).

20


**25 11E. 4-(tert-Butoxycarbonylamino-methyl)-3-fluorobenzoic acid methyl ester**

To a solution of 4-aminomethyl-3-fluorobenzoic acid methyl ester (1.5 g, 8.2 mmol) in dichloromethane (20 ml) were added di-*tert*-butyl dicarbonate (2.3 g, 11 mmol) and triethylamine (1.4 ml, 10 mmol). The mixture was stirred for 18h, washed with 0.3M KHSO<sub>4</sub> and brine and concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant 10% EtOAc/pet. ether) to give a white solid identified as 4-(tert-butoxycarbonylamino-methyl)-3-fluorobenzoic acid methyl ester (1.4 g, 60%),

30

**11F. 4-(tert-Butoxycarbonylamino-methyl)-3-fluorobenzoic acid**


To a solution of 4-(tert-Butoxycarbonylamino-methyl)-3-fluorobenzoic acid methyl ester (640 mg, 2.25 mmol) in dioxan (40 ml) was added 1N NaOH (4.5 ml, 4.5 mmol). The mixture was stirred for 18 h, diluted with ethyl acetate, washed with 1N KHSO<sub>4</sub>, water and brine and concentrated *in vacuo* to give a white solid identified as 4-(tert-butoxycarbonylamino-methyl)-3-fluorobenzoic acid (608 mg, 100%).

**Example 12****10 4-Cyano-3-ethylbenzoic acid****12A. 4-Bromo-2-ethylbenzonitrile**

4-Bromo-2-ethylaniline (12.5 g, 62.5 mmol) was reacted following the method of Example 9A to give a pale brown oil identified as 4-bromo-2-ethylbenzonitrile (7.66 g, 58%).

**12B. 4-Cyano-3-ethylbenzoic acid**

4-Bromo-2-ethylbenzonitrile (7.55 g, 35.9 mmol) was reacted following the method of Example 6 to give a pale brown solid identified as 4-cyano-3-ethylbenzoic acid (4.34 g, 69%).

**Example 13****4-Cyano-2-methoxybenzoic acid**

**13A. 4-Cyano-2-methoxybenzoic acid methyl ester**

4-Amino-2-methoxybenzoic acid methyl ester (4.59 g, 25.33 mmol) was reacted following the method of Example 9A to give a pale yellow solid identified as 4-cyano-2-methoxybenzoic acid methyl ester (2.58 g, 53%).

5


**13B. 4-Cyano-2-methoxybenzoic acid**

4-Cyano-2-methoxybenzoic acid methyl ester (2.68 g, 15.52 mmol) was reacted following the method of Example 10C to give a white powder identified as 4-cyano-2-methoxybenzoic acid (2.60 g, 95%).

10

**Example 14**

**4-(3-Methyl-4-(piperazine-1-carboxylaminomethyl)benzoyl)-5,6,7,8-tetrahydrothieno[3,2-*b*]azepine hydrochloride**



15

**14A: 4-(3-Methyl-4-cyanobenzoyl)-5,6,7,8-tetrahydrothieno[3,2-*b*]azepine**

Thionyl chloride (5 ml, 68.55 mmol) was added to a stirred suspension of 4-cyano-3-methylbenzoic acid (1.43 g, 8.90 mmol) in dichloromethane (20 ml). The mixture was heated at reflux for 2 h, cooled to room temperature and concentrated *in vacuo*. The residue was azeotroped with dichloromethane then dissolved in dichloromethane 20 ml. The resulting solution was slowly added to a stirred solution of 5,6,7,8-tetrahydrothieno[3,2-*b*]azepine (1.36 g, 8.90 mmol) and triethylamine (3.70 ml, 26.54 mmol) in dichloromethane (30 ml). The mixture was stirred at room temperature for 24 h, washed with 1M

25 KHSO<sub>4</sub>, saturated NaHCO<sub>3</sub> and brine, then concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant 25% EtOAc/pet. ether) to give a brown solid identified as 4-(3-methyl-4-cyanobenzoyl)-5,6,7,8-tetrahydrothieno[3,2-*b*]azepine (1.70 g, 71%).

**14B: 4-(4-Aminomethyl-3-methylbenzoyl)-5,6,7,8-tetrahydrothieno[3,2-*b*]azepine**

Cobalt(II) chloride hexahydrate (2.84 g, 11.94 mmol) was added to a solution of 4-(3-methyl-4-cyanobenzoyl)-5,6,7,8-tetrahydrothieno[3,2-*b*]azepine (1.70 g, 5.70 mmol) in methanol (70 ml) at 0°C. Sodium borohydride (2.22 g, 58.68 mmol) was added portion wise at 0°C and the mixture was stirred at 0°C for 30 min then at room temperature for 2 h. Saturated ammonium chloride was then added and the mixture was stirred for 30 min then concentrated *in vacuo*. The residue was azeotroped with toluene then extracted with chloroform. The extracts were washed with brine and concentrated *in vacuo* to give a white solid identified as 4-(4-aminomethyl-3-methylbenzoyl)-5,6,7,8-tetrahydrothieno[3,2-*b*]azepine (1.12 g, 65%).

**14C: 4-(4-(4-(*tert*-Butyloxycarbonyl)piperazine-1-carboxylaminomethyl)-3-methylbenzoyl)-5,6,7,8-tetrahydrothieno[3,2-*b*]azepine**

1,1'-Carbonyldiimidazole (234 mg, 1.45 mmol) was added to a solution of 4-(4-aminomethyl-3-methylbenzoyl)-5,6,7,8-tetrahydrothieno[3,2-*b*]azepine (400 mg, 1.33 mmol) and DIEA (0.3 ml, 1.72 mmol) in DMF (20 ml) and the mixture was stirred at room temperature for 30 min. *tert*-Butyl piperazine-1-carboxylate (281 mg, 1.50 mmol) was added and the mixture was stirred at room temperature for 24 h then concentrated *in vacuo*. The residue was taken up in chloroform and the solution was washed with 1M KHSO<sub>4</sub> and brine, then concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant 75% EtOAc/pet. ether) to give a white solid identified as 4-(4-(4-(*tert*-butyloxycarbonyl)piperazine-1-carboxylaminomethyl)-3-methylbenzoyl)-5,6,7,8-tetrahydrothieno[3,2-*b*]azepine (588 mg, 86%).

**14D: 4-(3-Methyl-4-(piperazine-1-carboxylaminomethyl)benzoyl)-5,6,7,8-tetrahydrothieno[3,2-*b*]azepine hydrochloride**


A solution of 4-(4-(4-(*tert*-butyloxycarbonyl)piperazine-1-carboxylaminomethyl)-3-methylbenzoyl)-5,6,7,8-tetrahydrothieno[3,2-*b*]azepine (588 mg, 1.15 mmol) in 4N HCl/dioxan (10 ml) was stirred at room temperature for 30 min then concentrated *in vacuo*. The residue was dissolved in acetonitrile/water and lyophilised to give a white solid identified as 4-(3-methyl-4-(piperazine-1-carboxylaminomethyl)benzoyl)-5,6,7,8-tetrahydrothieno[3,2-*b*]azepine hydrochloride (393 mg, 76%).

<sup>1</sup>H NMR: d<sub>6</sub>-DMSO δ 1.60-1.74 (2H, m), 1.82-1.94 (2H, m), 2.17 (3H, s), 2.86-2.95 (2H, m), 2.96-3.10 (4H, m), 3.35-3.45 (2H, m), 3.50-3.64 (4H, m), 4.16 (2H, s), 6.26 (1H, br s); 6.85-7.10 (4H, m), 7.24 (1H, br s), 9.28 (1H, br s) ppm.

MS: [M+H]<sup>+</sup> = 413.2

### Example 15

10 5-(4-(4-Cyclopropylmethylpiperazine-1-carbonylaminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine



15A: 5-(4-Cyano-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*]-  
15 [1,5]benzodiazepine

Thionyl chloride (1.8 ml, 27 mmol) was added to a stirred suspension of 4-cyano-3-methylbenzoic acid (1.29 g, 8.0 mmol) in toluene (25 ml). The mixture was heated at reflux for 2 h, cooled to room temperature and concentrated *in vacuo*. The residue was azeotroped with toluene then dissolved in dichloromethane (10 ml). The resulting solution was added to a stirred suspension of 1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine (1.6 g, 8 mmol) and triethylamine (1.4 ml, 10 mmol) in dichloromethane (15 ml). The mixture was stirred overnight at room temperature then concentrated *in vacuo*. The residue was partitioned between chloroform and 0.3M KHSO<sub>4</sub>. The aqueous phase was extracted with chloroform/2-propanol (80:20). The combined organic phases were washed with sat. NaHCO<sub>3</sub> and brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant 5% methanol/chloroform) to give a pale yellow solid identified as 5-(4-cyano-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]-  
30 benzodiazepine (2.4 g, 87%).

**15B: 5-(4-Aminomethyl-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine**

Cobalt(II) chloride hexahydrate (1.59 g, 6.7 mmol) was added to an ice-cold solution of 5-(4-cyano-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine (1.15 g, 3.35 mmol) in methanol (35 ml). Sodium borohydride (1.27 g, 33.5 mmol) was added portion wise at 0°C and the mixture was stirred at RT for 1 h, then quenched with 1M KHSO<sub>4</sub> and concentrated *in vacuo*. The aqueous residue was diluted with 1M KHSO<sub>4</sub> (40ml) and filtered through Celite® filter agent. The filtrate was washed with diethyl ether (2 × 50 ml) then basified with 2 M NaOH and extracted with chloroform. The organic phase was dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated *in vacuo* to give a pale brown solid identified as 5-(4-aminomethyl-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine (745 mg, 64%).

**15C: 5-(4-(4-(*tert*-Butyloxycarbonyl)piperazine-1-carbonylamino-methyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]-benzodiazepine**

1,1'-Carbonyldiimidazole (76 mg, 0.47 mmol) was added to a solution of 5-(4-(aminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine (150 mg, 0.43 mmol) and DIEA (0.1 ml, 0.57 mmol) in DMF (10 ml). The solution was stirred for 30 min, *tert*-butyl piperazine-1-carboxylate (91 mg, 0.49 mmol) was added and stirring was continued for 72 h. The mixture was concentrated *in vacuo* and the residue was taken up in chloroform. The solution was washed with water and brine, dried and concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant 100% EtOAc then 10% methanol/EtOAc) to give a white solid identified as 5-(4-(4-(*tert*-butyloxycarbonyl)piperazine-1-carbonylamino-methyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]-benzodiazepine (160 mg, 66%).

**15D: 1-Methyl-5-(3-methyl-4-(piperazine-1-carbonylamino-methyl)-benzoyl)-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine hydrochloride**

A solution of 5-(4-(4-(*tert*-butyloxycarbonyl)piperazine-1-carbonylamino-methyl)-3-methylbenzoyl)-1-methyl-4,10-

dihydropyrazolo[5,4-*b*][1,5]benzodiazepine (160 mg, 0.29 mmol) in 4N HCl/dioxan (15 ml) was stirred at room temperature for 30 min then concentrated *in vacuo*. The residue was azeotroped with diethyl ether to give a white solid identified as 1-methyl-5-(3-methyl-4-(piperazine-1-carboxylaminomethyl)-benzoyl)-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine hydrochloride (130 mg, 90%).

**15E: 5-(4-(4-Cyclopropylmethylpiperazine-1-carboxylaminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine**

To a solution of 1-methyl-5-(3-methyl-4-(piperazine-1-carboxylaminomethyl)-benzoyl)-4,10-dihydropyrazolo[5,4-*b*][1,5]-benzodiazepine hydrochloride (100 mg, 0.20 mmol) and triethylamine (0.5 ml, 3.59 mmol) in THF (10 ml) were added cyclopropanecarboxaldehyde (14 mg, 0.20 mmol) and sodium cyanoborohydride (15 mg, 0.24 mmol) and the resulting mixture was stirred at room temperature for 24 h then concentrated *in vacuo*. The residue was dissolved in ethyl acetate and the resulting solution was washed with saturated NaHCO<sub>3</sub>, water and brine, dried and concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant 10% methanol/EtOAc) to give a white solid identified as 5-(4-(4-cyclopropylmethylpiperazine-1-carboxylaminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine (35 mg, 35%).


<sup>1</sup>H NMR: d<sub>4</sub>-MeOH δ 0.14 (2H, q, J=4.7Hz), 0.51-0.59 (2H, m), 0.82-0.95 (1H, m), 2.15 (3H, s), 2.28 (2H, d, J=6.7Hz), 2.52 (4H, t, J=4.9Hz), 3.43 (4H, t, J=4.9Hz), 3.80 (3H, s), 3.95 (1H, d, J=14.4Hz), 4.23 (2H, s), 5.78 (1H, d, J=14.6Hz), 6.61-6.74 (2H, m), 6.99 (2H, s), 7.03 (1H, s), 7.05-7.14 (1H, m), 7.19-7.24 (2H, m) ppm.

MS: [M+H]<sup>+</sup> = 514.3



## Example 16

5-(4-(4-Benzylpiperazine-1-carboxylaminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine



5

To a solution of 1-methyl-5-(3-methyl-4-(piperazine-1-carboxylaminomethyl)-benzoyl)-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine hydrochloride (100 mg, 0.20 mmol) and triethylamine (0.5 ml, 3.59 mmol) in THF (10 ml) were added benzaldehyde (21 mg, 0.20 mmol) and sodium cyanoborohydride (15 mg, 0.24 mmol) and the resulting mixture was stirred at room temperature for 24 h then concentrated *in vacuo*. The residue was dissolved in ethyl acetate and the resulting solution was washed with saturated NaHCO<sub>3</sub>, water and brine, dried and concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant 5% methanol/EtOAc) to give a white solid identified as 5-(4-(4-benzylpiperazine-1-carboxylaminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine (37 mg, 34%).

15

<sup>1</sup>H NMR: δ 2.10 (3H, s), 2.36-2.48 (4H, m), 3.29-3.44 (4H, m), 3.48-3.51 (2H, m), 3.76 (3H, s), 3.96 (1H, d, J=14.6Hz), 4.22-4.28 (2H, m), 4.61-4.68 (1H, m), 5.88 (1H, d, J=14.6Hz), 6.46 (1H, s), 6.62-6.74 (2H, m), 6.82-6.96 (3H, m), 6.98-7.11 (2H, m), 7.19-7.34 (5H, m) ppm.


20

MS: [M+H]<sup>+</sup> = 550.2

25

**Example 17**

**5-(4-(4-(3-Hydroxybenzyl)piperazine-1-carboxylaminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine**



5

**17A: 3-(*tert*-Butyldimethylsilyloxy)toluene**

*tert*-Butyldimethylsilyl chloride (3.00 g, 22.00 mmol) was added to a solution of *m*-cresol (2.00g, 18.00mmol) and triethylamine (4 ml, 28.7 mmol) in dichloromethane (50 ml) at 0°C. The mixture was stirred at room temperature for 24h then concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant 10% EtOAc/pet. ether) to give a colourless oil identified as 3-(*tert*-butyldimethylsilyloxy)toluene (3.60 g, 88%).

10

**17B: 3-(*tert*-Butyldimethylsilyloxy)benzyl bromide**

*N*-Bromosuccinimide (2.90 g, 16.20 mmol) and AIBN (266 mg, 1.62 mmol) were added to a stirred solution of 3-(*tert*-butyldimethylsilyloxy)toluene (3.60 g, 16.20 mmol) in carbon tetrachloride (120 ml) and the mixture was heated at reflux for 24h, then allowed to cool to room temperature and concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant cyclohexane) to give a colourless oil identified as 3-(*tert*-butyldimethylsilyloxy)benzyl bromide (2.45 g, 50%).

15

20

**17C: *tert*-Butyl 4-(3-hydroxybenzyl)piperazine-1-carboxylate**

Sodium hydride (406 mg, 60% dispersion in oil, 10.15 mmol) was added portionwise to a stirred solution of *tert*-butyl piperazine-1-carboxylate in DMF (50 ml) at 0°C. The mixture was allowed to warm to room temperature over 1h, then a solution of 3-(*tert*-butyldimethylsilyloxy)benzyl bromide (2.44 g, 8.10 mmol) in DMF (10 ml) was added dropwise and the mixture was stirred at room temperature for 24 h. Water was added and the mixture was stirred for 30min then poured into EtOAc. The organic phase was washed with saturated

25

30

NaHCO<sub>3</sub> and brine, then concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant 40% EtOAc/pet. ether) to give a light brown oil identified as *tert*-butyl 4-(3-hydroxybenzyl)piperazine-1-carboxylate (2.00 g, 84%).

5

**17D: 1-(3-Hydroxybenzyl)piperazine dihydrochloride**

A solution of *tert*-butyl 4-(3-hydroxybenzyl)piperazine-1-carboxylate (1.94 g, 6.60 mmol) in 4N HCl/dioxan (10ml) was stirred at room temperature for 30 min then concentrated *in vacuo*. The residue was triturated with diethyl  
10 ether to give a white solid identified as 1-(3-hydroxybenzyl)piperazine dihydrochloride (1.10 g, 63%).


**17E:** 5-(4-(4-(3-Hydroxybenzyl)piperazine-1-carboxylaminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine  
15 1,1'-Carbonyldiimidazole (15 mg, 0.09 mmol) was added to a stirred solution of 5-(4-(aminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine (31 mg, 0.09 mmol) and DIEA (0.1 ml 0.57 mmol) in DMF (5 ml). The solution was stirred for 1h, 1-(3-hydroxybenzyl)piperazine dihydrochloride (27 mg, 0.10 mmol) was added and  
20 stirring was continued at room temperature for 24 h. The mixture was concentrated *in vacuo* and the residue was taken up in EtOAc. The solution was washed with saturated NaHCO<sub>3</sub> and brine, then concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant 20% methanol/EtOAc) to give a white solid identified as 5-(4-(4-(3-hydroxybenzyl)-  
25 piperazine-1-carboxylaminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine (45 mg, 90%).

<sup>1</sup>H NMR: δ 2.15 (3H, s), 2.41 (4H, t, J=4.7Hz), 3.40 (4H, t, J=4.7Hz), 3.46 (2H, s), 3.80 (3H, s), 3.97 (1H, d, J=14.6Hz), 4.22 (2H, s), 4.90 (1H, m), 5.78  
30 (1H, d, J=14.6Hz), 6.62-6.79 (5H, m), 6.99 (2H, s), 7.03-7.27 (6H, m) ppm.

MS: [M+H]<sup>+</sup> = 566.1

**Example 18**

**5-(4-(4-(3-Hydroxymethylbenzyl)piperazine-1-carboxylaminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine**



5

**18A: *tert*-Butyl 4-(3-(methyloxycarbonyl)benzyl)piperazine-1-carboxylate**

Methyl 3-(bromomethylbenzoate) (1.23 g, 5.37 mmol) was added to a stirred solution of *tert*-butyl piperazine-1-carboxylate (1.00 g, 5.37 mmol) and triethylamine (1.50 ml, 10.74 mmol) in dichloromethane (20 ml). The solution was stirred at room temperature for 24h then concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant EtOAc) to give a white solid identified as *tert*-butyl 4-(3-(methyloxycarbonyl)benzyl)-piperazine-1-carboxylate (1.55 g, 86%).

**18B: *tert*-Butyl 4-(3-carboxybenzyl)piperazine-1-carboxylate**

Lithium hydroxide monohydrate (339 mg, 9.27 mmol) was added to a solution of *tert*-butyl 4-(3-(methyloxycarbonyl)benzyl)piperazine-1-carboxylate (1.55 g, 4.63 mmol) in THF (10 ml) and water (2 ml). The solution was stirred at room temperature for 24 h then acidified to pH 5 with 0.3 M KHSO<sub>4</sub> and extracted successively with chloroform and dichloromethane. The combined extracts were concentrated *in vacuo* to give a white solid identified as *tert*-butyl 4-(3-carboxybenzyl)piperazine-1-carboxylate (1.09 g, 74%).

**18C: *tert*-Butyl 4-(3-(hydroxymethyl)benzyl)piperazine-1-carboxylate**

Isobutyl chloroformate (0.47 ml, 3.64 mmol) was slowly added to an ice-cold solution of *tert*-butyl 4-(3-carboxybenzyl)piperazine-1-carboxylate (1.06 g, 3.31 mmol) and *N*-methylmorpholine (0.80 ml, 7.28 mmol) in THF (15 ml). The solution was stirred at 0°C for 45 min and then filtered. The filtrate was added to an ice-cold solution of sodium borohydride (313 mg, 8.27 mmol) in water (10 ml). The stirred mixture was allowed to warm to room

temperature over 2 h and then concentrated *in vacuo*. The residue was taken up in EtOAc and the solution was washed with water and brine then concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant EtOAc) to give a white solid identified as *tert*-butyl 4-(3-(hydroxymethyl)benzyl)piperazine-1-carboxylate (230 mg, 23%).

**18D: 1-(3-(Hydroxymethyl)benzyl)piperazine dihydrochloride**

A solution of *tert*-butyl 4-(3-(hydroxymethyl)benzyl)piperazine-1-carboxylate (230mg, 0.75mmol) in 4N HCl/dioxan (10ml) was stirred at room temperature for 45min then concentrated *in vacuo*. The residue was azeotroped with toluene to give a white solid identified as 1-(3-(hydroxymethyl)benzyl)-piperazine dihydrochloride (158 mg, 75%).

**18E: 5-(4-(4-(3-Hydroxymethylbenzyl)piperazine-1-carbonylamino-methyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]-benzodiazepine**


1,1'-Carbonyldiimidazole (20 mg, 0.12 mmol) was added to a solution of 5-(4-(aminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo-[5,4-*b*][1,5]benzodiazepine (35 mg, 0.10 mmol) in DMF (3 ml). The solution was stirred for 1 h, a solution of 1-(3-(hydroxymethyl)benzyl)piperazine dihydrochloride (31 mg, 0.11 mmol) and DIEA (54  $\mu$ l, 0.30 mmol) in DMF (2 ml) was added and the mixture was stirred at room temperature for 24 h then concentrated *in vacuo*. The residue taken up in chloroform and the solution was washed with brine and concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant 7% methanol/chloroform) to give a white solid identified as 5-(4-(4-(3-hydroxymethylbenzyl)piperazine-1-carbonylamino-methyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo-[5,4-*b*][1,5]benzodiazepine (27 mg, 50%).

<sup>1</sup>H NMR:  $\delta$  2.00 (3H, s), 2.32-2.36 (4H, m), 3.32-3.45 (4H, m), 3.46 (2H, s), 3.63 (3H, s), 3.91 (1H, d, J=14.6Hz), 4.10-4.20 (1H, m), 4.66 (2H, s), 5.28-5.29 (1H, m), 5.80 (1H, d, J=14.3Hz), 6.50-7.30 (15H, m) ppm.

MS: [M+H]<sup>+</sup> = 580.3

## Example 19.

1-Methyl-5-(3-methyl-4-(4-(4-picolyl)piperazine-1-carboxylaminomethyl)-benzoyl)-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine



5

To a solution of 1-methyl-5-(3-methyl-4-(piperazine-1-carboxylaminomethyl)-benzoyl)-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine hydrochloride (100 mg, 0.20 mmol) and triethylamine (0.5 ml, 3.59 mmol) in THF (10 ml) were added 4-pyridinecarboxaldehyde (21 mg, 0.20 mmol) and sodium cyanoborohydride (15 mg, 0.24 mmol) and the resulting mixture was stirred at room temperature for 24 h then concentrated *in vacuo*. The residue was dissolved in ethyl acetate and the resulting solution was washed with saturated NaHCO<sub>3</sub>, water and brine, dried and concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant 10%-30% methanol/EtOAc) to give a white solid identified as 1-methyl-5-(3-methyl-4-(4-(4-picolyl)piperazine-1-carboxylaminomethyl)benzoyl)-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine (33 mg, 30%).

15

<sup>1</sup>H NMR: δ 2.13 (3H, s), 2.34-2.49 (4H, m), 3.29-3.47 (4H, m), 3.76 (3H, s), 3.96 (1H, d, J=14.8Hz), 4.25-4.27 (2H, d, J=4.7Hz), 4.50-4.60 (1H, m), 5.90 (1H, d, J=14.4Hz), 6.25 (1H, s), 6.63-6.71 (2H, m), 6.84 (2H, s), 6.92 (1H, s), 7.00-7.12 (2H, m), 7.25 (5H, s), 8.53 (2H, d, J=5.9Hz) ppm.


20

MS: [M+H]<sup>+</sup> = 551.1

25

**Example 20**

**5-(4-(4-(2-Hydroxyethyl)piperazine-1-carboxylaminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine**




5  
1,1'-Carbonyldiimidazole (20 mg, 0.19 mmol) was added to a solution of 5-(4-(aminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo-[5,4-*b*][1,5]benzodiazepine (31 mg, 0.09 mmol) in DMF (3 ml). The solution was stirred at room temperature for 1h, a solution of 1-(2-hydroxyethyl)-  
10 piperazine (13 mg, 0.10 mmol) in DMF (2 ml) was added and stirring was continued for 72 h. The solution was concentrated *in vacuo* and the residue was partitioned between chloroform and brine. The organic layer was separated and concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant 7% methanol/chloroform) to give a white solid identified as  
15 5-(4-(4-(2-hydroxyethyl)piperazine-1-carboxylaminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine (22 mg, 48%).

<sup>1</sup>H NMR; δ 2.09 (3H, s), 2.42-2.59 (6H, m), 2.91-3.01 (1H, m), 3.33-3.62 (6H, m), 3.67 (3H, s), 3.93-3.98 (1H, m), 4.20-4.23 (2H, m), 5.00-5.03 (1H, m), 5.84-5.90 (1H, m), 6.64-7.25 (9H, m) ppm.  
20

MS: [M+H]<sup>+</sup> = 504.2

**Example 21**

**1-Methyl-5-(3-methyl-4-(4-(3-(methylthio)propyl)piperazine-1-carbonylaminomethyl)benzoyl)-4,10-dihydropyrazolo[5,4-*b*][1,5]-benzodiazepine**



5

To a solution of 1-methyl-5-(3-methyl-4-(piperazine-1-carbonylaminomethyl)-benzoyl)-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine hydrochloride (100 mg, 0.20 mmol) and triethylamine (0.5 ml, 3.59 mmol) in THF (10 ml) were added 3-(methylthio)propionaldehyde (21 mg, 0.20 mmol) and sodium cyanoborohydride (15 mg, 0.24 mmol) and the resulting mixture was stirred at room temperature for 24 h then concentrated *in vacuo*. The residue was dissolved in ethyl acetate and the resulting solution was washed with saturated NaHCO<sub>3</sub>, water and brine, dried and concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant 20% methanol/EtOAc) to give a white solid identified as 1-methyl-5-(3-methyl-4-(4-(3-(methylthio)propyl)piperazine-1-carbonylaminomethyl)benzoyl)-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine (41 mg, 38%).

<sup>1</sup>H NMR: δ 1.63-1.80 (3H, m), 2.04-2.12 (4H, m), 2.33-2.42 (6H, m), 2.48 (2H, t, J=6.7Hz), 3.29-3.39 (4H, m), 3.71 (3H, s), 3.93 (1H, d, J=14.4Hz), 4.12-4.30 (2H, m), 4.57-4.70 (1H, m), 5.85 (1H, d, J=14.6Hz), 6.44 (1H, s), 6.59-6.71 (2H, m), 6.83-6.88 (2H, m), 6.92-7.08 (2H, m), 7.14-7.27 (2H, m) ppm.


25

MS: [M+H]<sup>+</sup> = 548.0



**Example 22**

**5-(4-(4-(2-Aminoethyl)piperazine-1-carboxylaminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine dihydrochloride**



5

**22A: Benzyl 4-(2-hydroxyethyl)piperazine-1-carboxylate**

Benzyl chloroformate (3.40 ml, 24.00 mmol) was slowly added to an ice-cold stirred solution of 1-(2-hydroxyethyl)piperazine (2.60 g, 20.00 mmol) and DIEA (7.0 ml, 40.0 mmol) in dichloromethane (75 ml). The mixture was allowed to warm to room temperature and stirred for 24 h then concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant 6% methanol/chloroform) to give a colourless gum identified as benzyl 4-(2-hydroxyethyl)piperazine-1-carboxylate (4.80 g, 91%).

15

**22B: Benzyl 4-(2-bromoethyl)piperazine-1-carboxylate**

Carbon tetrabromide (7.23 g, 21.80 mmol) was added to an ice-cold stirred solution of benzyl 4-(2-hydroxyethyl)piperazine-1-carboxylate (4.80 g, 18.20 mmol) in dichloromethane (50 ml). The solution was stirred for 5 min, triphenylphosphine (5.95 g, 22.70 mmol) was added, and the mixture was allowed to warm to room temperature and stirred for 3 h. Silica gel was added and the solvent was removed *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant 50% EtOAc/pet. ether) to give a colourless gum identified as benzyl 4-(2-bromoethyl)piperazine-1-carboxylate (3.45 g, 58%).

25

**22C: Benzyl 4-(2-(*tert*-butoxycarbonylamino)ethyl)piperazine-1-carboxylate**

Benzyl 4-(2-bromoethyl)piperazine-1-carboxylate (3.45 g, 10.55 mmol) was added to an ice-cold saturated solution of ammonia in ethanol (60 ml). The

30

mixture was allowed to warm to room temperature and stirred for 4 h, then concentrated *in vacuo*. The residue was triturated with diethyl ether. The resultant solid was suspended in dichloromethane (75 ml) and triethylamine (2.25 ml, 16.00 mmol). The suspension was cooled to 0°C and di-*tert*-butyl dicarbonate (2.40 g, 11.00 mmol) was added. The mixture was allowed to warm to room temperature and stirred for 24 h then concentrated *in vacuo*. The residue was taken up in EtOAc. The solution was washed with saturated NaHCO<sub>3</sub> and brine, then concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant 3% methanol/chloroform) to give a yellow gum identified as benzyl 4-(2-(*tert*-butyloxycarbonylamino)ethyl)piperazine-1-carboxylate (2.60 g, 68%).

#### 22D: *tert*-Butyl 2-(1-piperazinyl)ethylcarbamate

Hydrogen was passed through a degassed solution of benzyl 4-(2-(*tert*-butyloxycarbonylamino)ethyl)piperazine-1-carboxylate (2.60 g, 7.16 mmol) in methanol (50 ml) containing 10% palladium on carbon (500 mg) for 2 h. The reaction mixture was filtered through Celite® and the filtrate was concentrated *in vacuo* to give a yellow gum identified as *tert*-butyl 2-(1-piperazinyl)ethylcarbamate (1.60 g, 97%).

#### 22E: 5-(4-(4-(2-(*tert*-Butyloxycarbonylaminoethyl)piperazine-1-carbonylaminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo-[5,4-*b*][1,5]benzodiazepine

1,1'-Carbonyldiimidazole (25 mg, 0.15 mmol) was added to a solution of 5-(4-(aminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo-[5,4-*b*][1,5]benzodiazepine (31 mg, 0.09 mmol) and DIEA (0.1 ml, 0.57 mmol) in DMF (5 ml). The solution was stirred for 1h, *tert*-butyl 2-(1-piperazinyl)ethylcarbamate (22 mg, 0.10 mmol) was added and stirring was continued at room temperature for 24 h. The mixture was concentrated *in vacuo* and the residue was taken up in EtOAc. The solution was washed with saturated NaHCO<sub>3</sub> and brine, then concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant 20% methanol/EtOAc) to give a white solid identified as 5-(4-(4-(2-(*tert*-butyloxycarbonylaminoethyl)piperazine-1-carbonylaminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo-[5,4-*b*][1,5]benzodiazepine (44 mg, 81%).

**22F: 5-(4-(4-(2-Aminoethyl)piperazine-1-carboxylaminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine dihydrochloride**


A solution of 5-(4-(4-(2-(*tert*-butoxycarbonylaminoethyl)piperazine-1-carboxylaminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo-  
5 [5,4-*b*][1,5]benzodiazepine (42 mg, 0.07 mmol) in 4N HCl/dioxan (5 ml) was stirred at room temperature for 30 min then concentrated *in vacuo*. The residue was dissolved in acetonitrile/water and lyophilised to give a white solid identified as 5-(4-(4-(2-aminoethyl)piperazine-1-carboxylaminomethyl)-3-methyl-  
10 benzoyl)-1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine dihydrochloride (37 mg, 92%).

$^1\text{H}$  NMR:  $\delta$  2.17 (3H, s), 3.30-3.35 (4H, m), 3.41-3.50 (1H, m), 3.56-3.72 (4H, m), 4.00 (3H, s), 4.04 (1H, s), 4.26 (2H, s), 4.83-4.89 (2H, m), 5.88 (1H, d,  $J=15\text{Hz}$ ), 6.83-6.84 (2H, m), 6.92-7.13 (4H, m), 7.15-7.28 (1H, m), 7.36 (1H, d,  $J=7.9\text{Hz}$ ), 7.96 (1H, s) ppm.

MS:  $[\text{M}+\text{H}]^+ = 503.5$

**20 Example 23**

**1-Methyl-5-(3-methyl-4-(4-methylperhydro-1,4-diazepine-1-carboxylaminomethyl)benzoyl)-4,10-dihydropyrazolo[5,4-*b*][1,5]-benzodiazepine**



25

1,1'-Carbonyldiimidazole (37 mg, 0.23 mmol) was added to a solution of 5-(4-(aminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo-  
[5,4-*b*][1,5]benzodiazepine (75 mg, 0.22 mmol) in DMF (2 ml). The solution was stirred for 1h, a solution of 1-methylhomopiperazine (27 mg, 0.24 mmol) and DIEA (31 mg, 0.24 mmol) in DMF (1 ml) was added and stirring was con-  
30


tinued for 24 h. The mixture was concentrated *in vacuo* and the residue was purified by chromatography on silica gel (eluant 30/2/1 – 1/1/1 chloroform/methanol/concentrated ammonia) to give a white solid identified as 1-methyl-5-(3-methyl-4-(4-methylperhydro-1,4-diazepine-1-carbonylamino-methyl)benzoyl)-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine (38 mg, 36%).

<sup>1</sup>H NMR: δ 1.80-1.99 (2H, m), 2.10 (3H, s), 2.35 (3H, s), 2.51-2.69 (4H, m), 3.39 (2H, t, J=5.9Hz), 3.45-3.68 (2H, m), 3.63 (3H, s), 3.95 (1H, d, J=14.6Hz), 4.23 (2H, t, J=4.2Hz), 4.65-4.75 (1H, m), 5.85 (1H, d, J=14.6Hz), 6.65-6.75 (2H, m), 6.76-6.88 (2H, m), 6.90-7.09 (2H, m), 7.11-7.22 (2H, m) ppm.

MS: [M+H]<sup>+</sup> = 488.2

#### Example 24

5-(4-(4-(2-Hydroxyethyl)piperazine-1-carbonylamino-methyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[4,5-*c*]pyrido[2,3-*b*][1,4]diazepine



24A: 5-(4-Cyano-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[4,5-*c*]pyrido[2,3-*b*][1,4]diazepine

Thionyl chloride (0.6 ml, 9.00 mmol) was added to a suspension of 4-cyano-3-methylbenzoic acid (322 mg, 2.00 mmol) in toluene (10 ml). The mixture was heated at reflux for 2 h, allowed to cool and concentrated *in vacuo*. The residue was azeotroped with toluene and then taken up in dichloromethane (5 ml). The solution was added slowly to a stirred solution of 1-methyl-4,10-dihydropyrazolo[4,5-*c*]pyrido[2,3-*b*][1,4]diazepine (400 mg, 2.00 mmol) and triethylamine (0.35 ml, 2.50 mmol) in dichloromethane (5 ml). The mixture was stirred at room temperature for 24 h then concentrated *in vacuo*. The resi-

due was purified by flash chromatography on silica gel (eluant 5% methanol/chloroform) to give an orange solid identified as 5-(4-cyano-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[4,5-*c*]pyrido[2,3-*b*][1,4]diazepine (500 mg, 73%).

5

**24B: 5-(4-Aminomethyl-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[4,5-*c*]pyrido[2,3-*b*][1,4]diazepine**

Cobalt(II) chloride hexahydrate (690 mg, 2.90 mmol) was added to an ice-cold stirred solution of 5-(4-cyano-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[4,5-*c*]pyrido[2,3-*b*][1,4]diazepine (500 mg, 1.45 mmol) in methanol (15 ml). Sodium borohydride (570 mg, 15.00 mmol) was added portion wise and the mixture was stirred at room temperature for 1h. 1M KHSO<sub>4</sub> was added, the methanol was removed *in vacuo*, and the aqueous residue was filtered through Celite®. The filtrate was washed with diethyl ether, basified to pH12 with 2M sodium hydroxide and extracted with chloroform. The chloroform extracts were washed with brine and concentrated *in vacuo* to give a pale orange solid identified as 5-(4-aminomethyl-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[4,5-*c*]pyrido[2,3-*b*][1,4]diazepine (400 mg, 79%).

**24C: 5-(4-(4-(2-Hydroxyethyl)piperazine-1-carbonylaminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[4,5-*c*]pyrido[2,3-*b*][1,4]diazepine**

1,1'-Carbonyldiimidazole (20 mg, 0.12 mmol) was added to a solution of 5-(4-aminomethyl-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[4,5-*c*]pyrido[2,3-*b*][1,4]diazepine (35 mg, 0.10 mmol) in DMF (3 ml). The solution was stirred for 1h, a solution of 1-(2-hydroxyethyl)piperazine (13 mg, 0.10 mmol) and DIEA (18 µl, 0.10 mmol) in DMF (2 ml) was added and the mixture was stirred at room temperature for 24 h then concentrated *in vacuo*. The residue taken up in chloroform and the solution was washed with brine and concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant 7% methanol/chloroform) to give a pale yellow solid identified as 5-(4-(4-(2-hydroxyethyl)piperazine-1-carbonylaminomethyl)-3-methylbenzoyl)-1-methyl-4,10-dihydropyrazolo[4,5-*c*]pyrido[2,3-*b*][1,4]diazepine (29 mg, 58%).


35

<sup>1</sup>H NMR:  $\delta$  2.42 (3H, br s), 2.44-2.60 (7H, m), 3.20-3.40 (4H, m), 3.55-3.65 (2H, m), 3.79 (3H, s), 3.85-4.00 (1H, m), 4.26 (2H, br s), 4.88 (1H, br s), 5.80-5.95 (1H, m), 6.60 (1H, br s), 6.80-7.30 (6H, m), 8.00 (1H, s) ppm.

5 MS:  $[M+H]^+ = 505.2$

### Example 25

4-(2-Cyclopropyl-ethyl)-piperazine-1-carboxylic acid 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide



25A: 3-Methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzonitrile

Thionyl chloride (0.8 ml, 11.00 mmol) was added to a suspension of 4-Cyano-2-methylbenzoic acid (630 mg, 3.91 mmol) in toluene (30 ml). The mixture was heated at reflux for 2h, allowed to cool and concentrated *in vacuo*. The residue was azeotroped with toluene and then taken up in dichloromethane (25ml). The resulting solution was added to a stirred solution of 1-methyl-4,10-dihydropyrazolo[5,4-*b*][1,5]benzodiazepine (706 mg, 3.53 mmol) and triethylamine (0.70ml, 5.02mmol) in dichloromethane (25ml). The mixture was heated at reflux for 18h and cooled. The mixture was diluted with dichloromethane and washed with 0.3 M KHSO<sub>4</sub>, saturated NaHCO<sub>3</sub> and brine then concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant ethyl acetate) to give an off-white solid identified as 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzonitrile (1.06 g, 87%).

**25B: (4-Aminomethyl-2-methyl-phenyl)-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulen-9-yl)-methanone**

Cobalt(II) chloride hexahydrate (1.45 g, 6.09 mmol) was added to a stirred solution of 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzonitrile (1.04 g, 3.03 mmol) in methanol (50 ml). Sodium borohydride (1.16 g, 30.66 mmol) was added portion-wise and the mixture was stirred at room temperature for 3 h. 0.3 M KHSO<sub>4</sub> was added and the mixture was concentrated *in vacuo*. The residue was diluted with saturated NaHCO<sub>3</sub> and extracted with chloroform. The chloroform extracts were washed with brine, concentrated *in vacuo* and freeze dried to give an off-white solid identified as (4-aminomethyl-2-methyl-phenyl)-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulen-9-yl)-methanone (475 mg, 45%).

**25C: 4-(2-Cyclopropyl-ethyl)-piperazine-1-carboxylic acid 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide**


1,1'-Carbonyldiimidazole (29 mg, 0.18 mmol) was added to a solution of (4-aminomethyl-2-methyl-phenyl)-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulen-9-yl)-methanone (60 mg, 0.17 mmol) in DMF (3 ml). The solution was stirred for 1 h, a solution of 1-(2-Cyclopropyl-ethyl)-piperazine (39 mg, 0.17 mmol) and DIEA (98 µl, 0.55 mmol) in DMF (1 ml) was added and the mixture was stirred at room temperature for 18 h. The mixture was diluted with ethyl acetate and washed with 5% KHCO<sub>3</sub>, water and brine and concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (eluant 100/10/1 chloroform/methanol/concentrated ammonia) to give a white solid identified as 4-(2-cyclopropyl-ethyl)-piperazine-1-carboxylic acid 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide (51 mg, 56%).

<sup>1</sup>H NMR: δ 0.01-0.09 (2H, m) 0.42-0.44 (2H, m), 0.62-0.66 (1H, m), 1.34-1.40 (2H, m), 2.29 (3H, s) 2.31-2.42 (6H, m), 3.31-3.35 (4H, m), 3.66 (3H, s), 3.97 (1H, d, J=14.6Hz), 4.20 (2H, d, J=5.4Hz), 4.39-4.41 (1H, m), 4.89-4.94 (1H, m), 5.83 (1H, d, J=14.6Hz) 6.56-7.26 (8H, m)

MS: [M+H]<sup>+</sup> = 528.7

**Examples 26 – 150**

The following compounds were prepared using methods analogous to those described above.

**5 Examples 26 - 35**


	a	R <sup>1</sup>	R <sup>4</sup>	MS: [M+H] <sup>+</sup>
26	1	Cl	CH <sub>3</sub>	447.3
27	2	Cl	CH <sub>3</sub>	461.3
28	1	Me	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	469.3
29	1	Me	CH <sub>2</sub> CH(CH <sub>3</sub> )CH <sub>2</sub> CH <sub>3</sub>	483.3
30	1	Me	CH <sub>2</sub> -	467.3
31	1	Cl		510.3
32	1	Me	CH <sub>2</sub> -	533.3
33	1	Me	CH <sub>2</sub> -	523.2
34	1	Me	CH <sub>2</sub> -	539.3







59

	a	R <sup>1</sup>	R <sup>4</sup>	MS: [M+H] <sup>+</sup>
35	1	Me	CH <sub>2</sub> CH <sub>2</sub> OH	457.2



## Examples 36 - 51




5

	A <sup>3</sup>	A <sup>6</sup>	A <sup>8</sup>	b	R <sup>4</sup>	MS: [M+H] <sup>+</sup>
36	CH	NH	N-Me	3	H	474.1
37	CH	NH	N-Me	3	CH <sub>2</sub> CH(CH <sub>3</sub> )CH <sub>2</sub> CH <sub>3</sub>	544.3
38	CH	NH	N-Me	3	CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>3</sub>	544.3
39	CH	NH	N-Me	3	CH <sub>2</sub> - 	528.3
40	CH	N-Me	N-Me	1		514.3
41	N	NH	N-Me	2	CH <sub>2</sub> - 	680.2
42	CH	N-Me	N-Me	2	CH <sub>2</sub> - 	594.3
43	CH	NH	N-Me	3	CH <sub>2</sub> - 	570.3
44	CH	NH	N-Me	2		556.3

60

	A <sup>3</sup>	A <sup>6</sup>	A <sup>8</sup>	b	R <sup>4</sup>	MS: [M+H] <sup>+</sup>
45	CH	NH	N-Me	3		600.3
46	CH	NH	N-Me	2		586.3
47	N	NH	N-Me	2	CH <sub>2</sub> CH <sub>2</sub> NH <sub>2</sub>	504.1
48	N	NH	N-Me	2	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> NH <sub>2</sub>	518.3
49	CH	NH	N-Me	2		571.4
50	N	NH	N-Me	2		572.3
51	CH	NH	N-CH <sub>2</sub> Ph	2	CH <sub>2</sub> CH <sub>2</sub> OH	580.2








## Examples 52 - 126












5

	R <sup>4</sup>	MS: [M+H] <sup>+</sup>
52	H	460.2
53	CH <sub>3</sub>	474.2
54	CH <sub>2</sub> CH <sub>3</sub>	488.2
55	CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	502.3
56	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	516.3
57	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	530.3










61

	R <sup>4</sup>	MS: [M+H] <sup>+</sup>
58	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	544.3
59	CH <sub>2</sub> CH <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>	530.3
60	CH <sub>2</sub> CH(CH <sub>3</sub> )CH <sub>2</sub> CH <sub>3</sub>	530.3
61	CH <sub>2</sub> CH(CH <sub>2</sub> CH <sub>3</sub> ) <sub>2</sub>	544.3
62	CH <sub>2</sub> CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>3</sub>	544.2
63	CH <sub>2</sub> - 	556.3
64	CH <sub>2</sub> CH=CH <sub>2</sub>	500.1
65	CH <sub>2</sub> - 	528.3
66	--- 	536.2
67	CH <sub>2</sub> - 	580.3
68	CH <sub>2</sub> - 	580.3
69	CH <sub>2</sub> - 	580.2
70	CH <sub>2</sub> - 	608.3







62








	R <sup>4</sup>	MS: [M+H] <sup>+</sup>
71		634.2
72		634.2
73		634.2
74		540.3
75		570.2
76		614.2
77		586.3
78		648.2
79		620.2

63










	R <sup>4</sup>	MS: [M+H] <sup>+</sup>
80		538.2
81		553.1
82		540.2
83		629.2
84		587.3
85		599.2
86		551.3
87		609.1
88		581.3

64

	R <sup>4</sup>	MS: [M+H] <sup>+</sup>
89		516.3
90		530.2
91		592.2
92	CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	546.3
93	CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H	532.1
94	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	560.2
95	CH <sub>2</sub> CH <sub>2</sub> CN	513.4
96	CH <sub>2</sub> CH <sub>2</sub> N <sub>3</sub>	529.2
97		571.2
98		572.2
99		546.3





	R <sup>4</sup>	MS: [M+H] <sup>+</sup>
100		608.3
101		609.3
102		557.3
103		560.3
104		572.2
105	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> NH <sub>2</sub>	517.3
106	CH <sub>2</sub> CH <sub>2</sub> N(CH <sub>2</sub> CH <sub>3</sub> ) <sub>2</sub>	559.3
107		573.2
108		557.3
109	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> OH	532.3

66


	$R^4$	MS: [M+H] <sup>+</sup>
110		530.2
111		536.9 [M+Na] <sup>+</sup>
112		532.3
113		594.3
114		536.2
115		534.3
116		548.1
117		576.3
118	$\text{CH}_2\text{CH}_2\text{OCH}_3$	518.2
119	$\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_3$	532.3
120	$\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2\text{OCH}_3$	562.3
121	$\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2\text{OCH}_2\text{CH}_3$	576.3
122		594.3





67


	R <sup>4</sup>	MS: [M+H] <sup>+</sup>
123		528.4
124		544.4
125		542.6
126		528.7

## Examples 127 - 129




	A <sup>8</sup>	R <sup>4</sup>	MS: [M+H] <sup>+</sup>
127	CH		550.3
128	CH		592.3
129	N	CH <sub>2</sub> CH <sub>2</sub> OH	489.0

## Examples 130 - 137




	A <sup>6</sup>	R <sup>4</sup>	MS: [M+H] <sup>+</sup>
130	NH		584.9 [M+Na] <sup>+</sup>
131	NH		513.3
132	NH	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> NH <sub>2</sub>	514.3
133	NH	CH <sub>2</sub> CH <sub>2</sub> NH <sub>2</sub>	500.3
134	O	CH <sub>2</sub> CH <sub>2</sub> OH	502.1
135	NH	CH <sub>2</sub> CH <sub>2</sub> OH	501.3
136	NH		
137	NH		533.3

## Examples 138 - 141












	R <sup>1</sup>	R <sup>4</sup>	MS: [M+H] <sup>+</sup>
138	F		518.5
139	F		532.4
140	Cl		534.2
141	Cl		548.2

## Examples 142 - 149




70

	R <sup>3</sup>	R <sup>4</sup>	MS: [M+H] <sup>+</sup>
142	CH <sub>3</sub>		514.3
143	CH <sub>3</sub>		528.3
144	F		518.3
145	F		532.1
146	Cl		534.5
147	Cl		548.6
148	CH <sub>2</sub> CH <sub>3</sub>		
149	OCH <sub>3</sub>		544.4

**Example 150**

71

	R <sup>4</sup>	MS: [M+H] <sup>+</sup>
150		514.3

**Example 151****In vitro Testing**

Compounds were assayed to determine their ability to inhibit the cellular consequences of AVP stimulation on intact cells. In the assay, the compounds of the invention cause significant inhibition of cellular activation at concentrations of 30  $\mu$ M or less. Preferred compounds cause significant inhibition at concentrations of 300 nM

**Example 152****Tablet for Oral Administration**

Tablets containing 100 mg of the compound of Example 15 as the active agent are prepared from the following:

Compound of Example 15	200.0 g
Corn starch	71.0 g
Hydroxypropylcellulose	18.0 g
Carboxymethylcellulose calcium	13.0 g
Magnesium stearate	3.0 g
Lactose	195.0 g
<i>Total</i>	<i>500.0 g</i>


The materials are blended and then pressed to give 2000 tablets of 250 mg, each containing 100 mg of the compound of Example 15.

The forgoing demonstrates that the compounds according to the present invention act as antagonists at the vasopressin V<sub>1a</sub> receptor and accordingly they may find utility as pharmaceutical agents for the treatment of conditions such as primary dysmenorrhoea, pre-term labour, hypertension, Raynaud's disease, brain oedema, motion sickness, small cell lung cancer, depression, anxiety, hyponatremia, liver cirrhosis and congestive heart failure.

The scope of the present invention is further defined in the following claims.

73  
CLAIMS


1. A compound according to general formula 1, or a pharmaceutically acceptable salt thereof



1


wherein:

-  $G^1$  is selected from a group according to general formula 2, a group according to general formula 3, a group according to general formula 4, a group according to general formula 5, a group according to general formula 6 and a group according to general formula 7;




2


3




4



5



6



7

- A<sup>1</sup> is selected from CH<sub>2</sub>, CH(OH), NH, N-alkyl, O and S;
- A<sup>2</sup> is selected from CH<sub>2</sub>, CH(OH), C(=O) and NH;
- A<sup>3</sup> is selected from S, NH, N-alkyl, -CH=CH- and -CH=N-;
- A<sup>4</sup> and A<sup>5</sup> are each selected from CH and N;
- 5 - A<sup>6</sup> is selected from CH<sub>2</sub>, NH, N-alkyl and O;
- A<sup>7</sup> and A<sup>11</sup> are selected from C and N;
- A<sup>8</sup> and A<sup>9</sup> are selected from CH, N, NH, N(CH<sub>2</sub>)<sub>d</sub>R<sup>7</sup> and S;
- A<sup>10</sup> is selected from -CH=CH-, CH, N, NH, N(CH<sub>2</sub>)<sub>d</sub>R<sup>7</sup> and S;
- A<sup>12</sup> and A<sup>13</sup> are selected from N and C;
- 10 - A<sup>14</sup>, A<sup>15</sup> and A<sup>16</sup> are selected from NH, N-CH<sub>3</sub>, S, N and CH;
- X<sup>1</sup> is selected from O and NH;
- R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> are each selected from H, alkyl, O-alkyl, F, Cl and Br;
- R<sup>4</sup> is selected from H, alkyl, alkenyl, alkynyl, optionally substituted phenyl, optionally substituted thienyl, optionally substituted furyl, optionally substituted pyridyl, optionally substituted pyrrolyl, optionally substituted pyrazolyl,
- 15 optionally substituted imidazolyl, optionally substituted oxazolyl, optionally substituted isoxazolyl, optionally substituted thiazolyl, optionally substituted isothiazolyl, -(CH<sub>2</sub>)<sub>e</sub>R<sup>8</sup>, -CH<sub>2</sub>-CH=CH-CH<sub>2</sub>-R<sup>8</sup>, -CH<sub>2</sub>-C≡C-CH<sub>2</sub>-R<sup>8</sup>, -(CH<sub>2</sub>)<sub>g</sub>-CH(OH)-(CH<sub>2</sub>)<sub>h</sub>-R<sup>8</sup>, -(CH<sub>2</sub>)<sub>i</sub>-O-(CH<sub>2</sub>)<sub>j</sub>-R<sup>8</sup> and  $\text{CH}_2-\triangle-\text{R}^8$  ;
- 20 - R<sup>5</sup> and R<sup>6</sup> are independently selected from alkyl, Ar and -(CH<sub>2</sub>)<sub>f</sub>-Ar;
- R<sup>7</sup> is selected from H, alkyl, optionally substituted phenyl, F, OH, O-alkyl, O-acyl, S-alkyl, NH<sub>2</sub>, NH-alkyl, N(alkyl)<sub>2</sub>, NH-acyl, N(alkyl)-acyl, CO<sub>2</sub>H, CO<sub>2</sub>-alkyl, CONH<sub>2</sub>, CONH-alkyl, CON(alkyl)<sub>2</sub>, CN and CF<sub>3</sub>;
- R<sup>8</sup> is selected from H, alkyl, alkenyl, alkynyl, acyl, optionally substituted phenyl, optionally substituted pyridyl, optionally substituted thienyl, optionally substituted furyl, optionally substituted pyrrolyl, optionally substituted pyrazolyl, optionally substituted imidazolyl, optionally substituted oxazolyl, optionally substituted isoxazolyl, optionally substituted thiazolyl, optionally substituted isothiazolyl, F, OH, hydroxyalkyl, O-alkyl, O-acyl, S-alkyl, NH<sub>2</sub>,
- 25 NH-alkyl, N(alkyl)<sub>2</sub>, 1-pyrrolidinyl, 1-piperidinyl, 4-morpholinyl, NH-acyl, N(alkyl)-acyl, N<sub>3</sub>, CO<sub>2</sub>H, CO<sub>2</sub>-alkyl, CONH<sub>2</sub>, CONH-alkyl, CON(alkyl)<sub>2</sub>, CN and CF<sub>3</sub>;
- 30 - Ar is selected from optionally substituted thienyl and optionally substituted phenyl;



- a is 1 or 2, b is 1, 2 or 3; c is 1 or 2, d is 1, 2 or 3; e is 1, 2, 3 or 4; f is 1, 2 or 3 and g, h, i and j are all independently 1 or 2;

provided that:

- not more than one of  $A^8$ ,  $A^9$  and  $A^{10}$  is  $NH$ ,  $N(CH_2)_dR^7$  or  $S$ ;
- 5 -  $A^7$  and  $A^{11}$  are not both simultaneously  $N$ ;
- neither  $A^7$  nor  $A^{11}$  is  $N$  if one of  $A^8$ ,  $A^9$  and  $A^{10}$  is  $NH$ ,  $N(CH_2)_dR^7$  or  $S$ ;
- if  $A^{10}$  is not  $-CH=CH-$  then one of  $A^8$ ,  $A^9$  and  $A^{10}$  is  $NH$ ,  $N(CH_2)_dR^7$  or  $S$  or one of  $A^7$  and  $A^{11}$  is  $N$ ;
- not more than one of  $A^{14}$ ,  $A^{15}$  and  $A^{16}$  is  $NH$ ,  $N-CH_3$  or  $S$ ;
- 10 -  $A^{12}$  and  $A^{13}$  are not both simultaneously  $N$ ;
- if one of  $A^{14}$ ,  $A^{15}$  and  $A^{16}$  is  $NH$ ,  $N-CH_3$  or  $S$  then  $A^{12}$  and  $A^{13}$  are both  $C$ ;
- and
- one of  $A^{14}$ ,  $A^{15}$  and  $A^{16}$  is  $NH$ ,  $N-CH_3$  or  $S$  or one of  $A^{12}$  and  $A^{13}$  is  $N$ ,
- wherein said compound is selected from the group consisting of:
- 15 - 4-(3,3-Dimethyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 4-(2-Cyclopropyl-ethyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 20 - 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 3-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 4-(2-Hydroxymethyl-cyclopropylmethyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 25 - 4-(3-Methyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 4-Cyclopentylmethyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 30 - 4-Cyclohexylmethyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 3-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 35 - 4-(3-Methyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;

- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 4-(2-Cyclopropyl-ethyl)-piperazine-1-carboxylic acid 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 5 - 4-Pentyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 4-Hexyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- (R)-4-(2-Methyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 10 - 4-(2-Ethyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 4-(2-Methyl-but-2-enyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 15 - 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 2-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 20 - 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 2-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 25 - 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 2-ethyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 2-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 2-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- 30 4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- and
- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-methoxy-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide.

- 35           2. A pharmaceutical composition comprising a compound according to claim 1 as an active agent.

3. A pharmaceutical composition according to Claim 2 formulated as a tablet or capsule for oral administration.

5 4. A pharmaceutical composition according to Claim 2 or 3 for treatment of primary dysmenorrhoea.


5. A pharmaceutical composition according to Claim 2 or 3 for treatment of male erectile dysfunction.

10

6. A pharmaceutical composition according to Claim 2 or 3 for treatment of pre-term labour, hypertension, Raynaud's disease, brain oedema, motion sickness, small cell lung cancer, depression, anxiety, hyponatremia, liver cirrhosis or congestive heart failure.

15


7. The use of a compound according to general formula 1, or a pharmaceutically acceptable salt thereof




1

20 wherein:


-  $G^1$  is selected from a group according to general formula 2, a group according to general formula 3, a group according to general formula 4, a group according to general formula 5, a group according to general formula 6 and a group according to general formula 7;



2




3




4

25


78



5




6



5


7

- A<sup>1</sup> is selected from CH<sub>2</sub>, CH(OH), NH, N-alkyl, O and S;
- A<sup>2</sup> is selected from CH<sub>2</sub>, CH(OH), C(=O) and NH;
- A<sup>3</sup> is selected from S, NH, N-alkyl, -CH=CH- and -CH=N-;
- A<sup>4</sup> and A<sup>5</sup> are each selected from CH and N;
- 10 - A<sup>6</sup> is selected from CH<sub>2</sub>, NH, N-alkyl and O;
- A<sup>7</sup> and A<sup>11</sup> are selected from C and N;
- A<sup>8</sup> and A<sup>9</sup> are selected from CH, N, NH, N(CH<sub>2</sub>)<sub>d</sub>R<sup>7</sup> and S;
- A<sup>10</sup> is selected from -CH=CH-, CH, N, NH, N(CH<sub>2</sub>)<sub>d</sub>R<sup>7</sup> and S;
- A<sup>12</sup> and A<sup>13</sup> are selected from N and C;
- 15 - A<sup>14</sup>, A<sup>15</sup> and A<sup>16</sup> are selected from NH, N-CH<sub>3</sub>, S, N and CH;
- X<sup>1</sup> is selected from O and NH;
- R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> are each selected from H, alkyl, O-alkyl, F, Cl and Br;
- R<sup>4</sup> is selected from H, alkyl, alkenyl, alkynyl, optionally substituted phenyl, optionally substituted thienyl, optionally substituted furyl, optionally substituted pyridyl, optionally substituted pyrrolyl, optionally substituted pyrazolyl, optionally substituted imidazolyl, optionally substituted oxazolyl, optionally substituted isoxazolyl, optionally substituted thiazolyl, optionally substituted isothiazolyl, -(CH<sub>2</sub>)<sub>e</sub>R<sup>8</sup>, -CH<sub>2</sub>-CH=CH-CH<sub>2</sub>-R<sup>8</sup>, -CH<sub>2</sub>-C≡C-CH<sub>2</sub>-R<sup>8</sup>, -(CH<sub>2</sub>)<sub>g</sub>-CH(OH)-(CH<sub>2</sub>)<sub>h</sub>-R<sup>8</sup>, -(CH<sub>2</sub>)<sub>i</sub>-O-(CH<sub>2</sub>)<sub>j</sub>-R<sup>8</sup> and CH<sub>2</sub>--R<sup>8</sup> ;
- 25 - R<sup>5</sup> and R<sup>6</sup> are independently selected from alkyl, Ar and -(CH<sub>2</sub>)<sub>f</sub>-Ar;

- R<sup>7</sup> is selected from H, alkyl, optionally substituted phenyl, F, OH, O-alkyl, O-acyl, S-alkyl, NH<sub>2</sub>, NH-alkyl, N(alkyl)<sub>2</sub>, NH-acyl, N(alkyl)-acyl, CO<sub>2</sub>H, CO<sub>2</sub>-alkyl, CONH<sub>2</sub>, CONH-alkyl, CON(alkyl)<sub>2</sub>, CN and CF<sub>3</sub>;
  - R<sup>8</sup> is selected from H, alkyl, alkenyl, alkynyl, acyl, optionally substituted phenyl, optionally substituted pyridyl, optionally substituted thienyl, optionally substituted furyl, optionally substituted pyrrolyl, optionally substituted pyrazolyl, optionally substituted imidazolyl, optionally substituted oxazolyl, optionally substituted isoxazolyl, optionally substituted thiazolyl, optionally substituted isothiazolyl, F, OH, hydroxyalkyl, O-alkyl, O-acyl, S-alkyl, NH<sub>2</sub>, NH-alkyl, N(alkyl)<sub>2</sub>, 1-pyrrolidinyl, 1-piperidinyl, 4-morpholinyl, NH-acyl, N(alkyl)-acyl, N<sub>3</sub>, CO<sub>2</sub>H, CO<sub>2</sub>-alkyl, CONH<sub>2</sub>, CONH-alkyl, CON(alkyl)<sub>2</sub>, CN and CF<sub>3</sub>;
  - Ar is selected from optionally substituted thienyl and optionally substituted phenyl;
  - a is 1 or 2, b is 1, 2 or 3; c is 1 or 2, d is 1, 2 or 3; e is 1, 2, 3 or 4; f is 1, 2 or 3 and g, h, i and j are all independently 1 or 2; provided that:
    - not more than one of A<sup>8</sup>, A<sup>9</sup> and A<sup>10</sup> is NH, N(CH<sub>2</sub>)<sub>d</sub>R<sup>7</sup> or S;
    - A<sup>7</sup> and A<sup>11</sup> are not both simultaneously N;
    - neither A<sup>7</sup> nor A<sup>11</sup> is N if one of A<sup>8</sup>, A<sup>9</sup> and A<sup>10</sup> is NH, N(CH<sub>2</sub>)<sub>d</sub>R<sup>7</sup> or S;
    - if A<sup>10</sup> is not -CH=CH- then one of A<sup>8</sup>, A<sup>9</sup> and A<sup>10</sup> is NH, N(CH<sub>2</sub>)<sub>d</sub>R<sup>7</sup> or S or one of A<sup>7</sup> and A<sup>11</sup> is N;
    - not more than one of A<sup>14</sup>, A<sup>15</sup> and A<sup>16</sup> is NH, N-CH<sub>3</sub> or S;
    - A<sup>12</sup> and A<sup>13</sup> are not both simultaneously N;
    - if one of A<sup>14</sup>, A<sup>15</sup> and A<sup>16</sup> is NH, N-CH<sub>3</sub> or S then A<sup>12</sup> and A<sup>13</sup> are both C; and
    - one of A<sup>14</sup>, A<sup>15</sup> and A<sup>16</sup> is NH, N-CH<sub>3</sub> or S or one of A<sup>12</sup> and A<sup>13</sup> is N,
- for the manufacture of a pharmaceutical composition for treatment of primary dysmenorrhoea.




30

8. The use of a compound according to general formula 1, or a pharmaceutically acceptable salt thereof



wherein:

- $G^1$  is selected from a group according to general formula 2, a group according to general formula 3, a group according to general formula 4, a group according to general formula 5, a group according to general formula 6 and a group according to general formula 7;



- $A^1$  is selected from  $CH_2$ ,  $CH(OH)$ ,  $NH$ ,  $N$ -alkyl,  $O$  and  $S$ ;  
 -  $A^2$  is selected from  $CH_2$ ,  $CH(OH)$ ,  $C(=O)$  and  $NH$ ;  
 -  $A^3$  is selected from  $S$ ,  $NH$ ,  $N$ -alkyl,  $-CH=CH-$  and  $-CH=N-$ ;  
 -  $A^4$  and  $A^5$  are each selected from  $CH$  and  $N$ ;

- A<sup>6</sup> is selected from CH<sub>2</sub>, NH, N-alkyl and O;
- A<sup>7</sup> and A<sup>11</sup> are selected from C and N;
- A<sup>8</sup> and A<sup>9</sup> are selected from CH, N, NH, N(CH<sub>2</sub>)<sub>d</sub>R<sup>7</sup> and S;
- A<sup>10</sup> is selected from -CH=CH-, CH, N, NH, N(CH<sub>2</sub>)<sub>d</sub>R<sup>7</sup> and S;
- 5 - A<sup>12</sup> and A<sup>13</sup> are selected from N and C;
- A<sup>14</sup>, A<sup>15</sup> and A<sup>16</sup> are selected from NH, N-CH<sub>3</sub>, S, N and CH;
- X<sup>1</sup> is selected from O and NH;
- R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> are each selected from H, alkyl, O-alkyl, F, Cl and Br;
- R<sup>4</sup> is selected from H, alkyl, alkenyl, alkynyl, optionally substituted phenyl,
- 10 optionally substituted thienyl, optionally substituted furyl, optionally substituted pyridyl, optionally substituted pyrrolyl, optionally substituted pyrazolyl, optionally substituted imidazolyl, optionally substituted oxazolyl, optionally substituted isoxazolyl, optionally substituted thiazolyl, optionally substituted isothiazolyl, -(CH<sub>2</sub>)<sub>e</sub>R<sup>8</sup>, -CH<sub>2</sub>-CH=CH-CH<sub>2</sub>-R<sup>8</sup>, -CH<sub>2</sub>-C≡C-CH<sub>2</sub>-R<sup>8</sup>, -(CH<sub>2</sub>)<sub>g</sub>-
- 15 CH(OH)-(CH<sub>2</sub>)<sub>h</sub>-R<sup>8</sup>, -(CH<sub>2</sub>)<sub>i</sub>-O-(CH<sub>2</sub>)<sub>j</sub>-R<sup>8</sup> and  $\text{CH}_2-\triangle-\text{R}^8$  ;
- R<sup>5</sup> and R<sup>6</sup> are independently selected from alkyl, Ar and -(CH<sub>2</sub>)<sub>f</sub>-Ar;
- R<sup>7</sup> is selected from H, alkyl, optionally substituted phenyl, F, OH, O-alkyl, O-acyl, S-alkyl, NH<sub>2</sub>, NH-alkyl, N(alkyl)<sub>2</sub>, NH-acyl, N(alkyl)-acyl, CO<sub>2</sub>H, CO<sub>2</sub>-alkyl, CONH<sub>2</sub>, CONH-alkyl, CON(alkyl)<sub>2</sub>, CN and CF<sub>3</sub>;
- 20 - R<sup>8</sup> is selected from H, alkyl, alkenyl, alkynyl, acyl, optionally substituted phenyl, optionally substituted pyridyl, optionally substituted thienyl, optionally substituted furyl, optionally substituted pyrrolyl, optionally substituted pyrazolyl, optionally substituted imidazolyl, optionally substituted oxazolyl, optionally substituted isoxazolyl, optionally substituted thiazolyl, optionally substituted isothiazolyl, F, OH, hydroxyalkyl, O-alkyl, O-acyl, S-alkyl, NH<sub>2</sub>,
- 25 NH-alkyl, N(alkyl)<sub>2</sub>, 1-pyrrolidinyl, 1-piperidinyl, 4-morpholinyl, NH-acyl, N(alkyl)-acyl, N<sub>3</sub>, CO<sub>2</sub>H, CO<sub>2</sub>-alkyl, CONH<sub>2</sub>, CONH-alkyl, CON(alkyl)<sub>2</sub>, CN and CF<sub>3</sub>;
- Ar is selected from optionally substituted thienyl and optionally substituted
- 30 phenyl;
- a is 1 or 2, b is 1, 2 or 3; c is 1 or 2, d is 1, 2 or 3; e is 1, 2, 3 or 4; f is 1, 2 or 3 and g, h, i and j are all independently 1 or 2;
- provided that:
- not more than one of A<sup>8</sup>, A<sup>9</sup> and A<sup>10</sup> is NH, N(CH<sub>2</sub>)<sub>d</sub>R<sup>7</sup> or S;
- 35 - A<sup>7</sup> and A<sup>11</sup> are not both simultaneously N;

82

- neither  $A^7$  nor  $A^{11}$  is N if one of  $A^8$ ,  $A^9$  and  $A^{10}$  is NH,  $N(CH_2)_dR^7$  or S;
  - if  $A^{10}$  is not  $-CH=CH-$  then one of  $A^8$ ,  $A^9$  and  $A^{10}$  is NH,  $N(CH_2)_dR^7$  or S or one of  $A^7$  and  $A^{11}$  is N;
  - not more than one of  $A^{14}$ ,  $A^{15}$  and  $A^{16}$  is NH,  $N-CH_3$  or S;
  - 5 -  $A^{12}$  and  $A^{13}$  are not both simultaneously N;
  - if one of  $A^{14}$ ,  $A^{15}$  and  $A^{16}$  is NH,  $N-CH_3$  or S then  $A^{12}$  and  $A^{13}$  are both C; and
  - one of  $A^{14}$ ,  $A^{15}$  and  $A^{16}$  is NH,  $N-CH_3$  or S or one of  $A^{12}$  and  $A^{13}$  is N,
- for the manufacture of a pharmaceutical composition for treatment of pre-term  
10 labour, hypertension, Raynaud's disease, brain oedema, motion sickness, small cell lung cancer, depression, anxiety, hyponatremia, liver cirrhosis or congestive heart failure.

9. The use according to Claim 7 or 8, wherein at least one of  $R^1$ ,  $R^2$  and  
15  $R^3$  is H and at least one is not H.

10. The use according to any one of Claims 7 – 9, wherein one of  $R^1$ ,  $R^2$   
and  $R^3$  is selected from an alkyl group, an O-alkyl group, F, Cl and Br and the  
others are H.

20

11. The use according to any one of the Claims 7 – 10, wherein  $X^1$  is  
NH.

12. The use according to any one of the Claims 7 – 11, wherein a is 1  
25 and b is 2.

13. The use according to any one of the Claims 7 – 12, wherein  $G^1$  is a  
group according to general formula 3.

30 14. The use according to Claim 13, wherein c is 2.

15. The use according to Claim 13 or 14, wherein  $A^1$  is  $CH_2$  and  $A^2$  is  
NH.

35 16. The use according to Claim 13 or 14, wherein  $A^1$  is NH or N-alkyl  
and  $A^2$  is  $C(=O)$ .



17. The use according to Claim 13 or 14, wherein  $A^3$  is S and  $A^4$  and  $A^5$  are both CH.

5 18. The use according to any of Claims 13 – 17, wherein  $A^3$  is –CH=CH– and  $A^4$  and  $A^5$  are both CH.

10 19. The use according to any of Claims 13 – 17, wherein  $A^3$  is –CH=N– and  $A^4$  and  $A^5$  are both CH.

20. The use according to any of Claims 13 – 17, wherein  $A^3$  is –CH=CH–,  $A^4$  is CH and  $A^5$  is N.

15 21. The use according to any of Claims 7 – 12, wherein  $G^1$  is a group according to general formula 6 or 7

22. The use according to Claim 21, wherein  $A^3$  is S and  $A^4$  and  $A^5$  are both CH.

20 23. The use according to Claim 21, wherein  $A^3$  is –CH=CH– and  $A^4$  and  $A^5$  are both CH.

25 24. The use according to Claim 21, wherein  $A^3$  is –CH=N– and  $A^4$  and  $A^5$  are both CH.

25 25. The use according to Claim 21, wherein  $A^3$  is –CH=CH–,  $A^4$  is CH and  $A^5$  is N.

30 26. The use according to any one to Claims 7 – 12, wherein  $G^1$  is a group according to general formula 4 or 6.

27. The use according to Claim 26, wherein  $A^6$  is NH.

35 28. The use according to Claim 26 or 27, wherein  $A^8$  is NH or N– $(CH_2)_d$ – $R^7$ .

29. The use according to Claim 28, wherein  $A^9$  is N and  $A^{10}$  is CH.

30. The use according to Claim 7 or 8, wherein one of  $R^1$ ,  $R^2$  and  $R^3$  is selected from an alkyl group, an O-alkyl group, F, Cl and Br and the others are H and  $X^1$  is NH.

31. The use according to any one of Claims 7 or 8, wherein one of  $R^1$ ,  $R^2$  and  $R^3$  is selected from an alkyl group, an O-alkyl group, F, Cl and Br and the others are H and  $X^1$  is NH, a is 1 and b is 2.

32. The use according to Claim 7 or 8, wherein  $G^1$  is a group according to general formula 6,  $A^4$ ,  $A^5$  and  $A^{10}$  are all CH,  $A^6$  is NH,  $A^7$  and  $A^{11}$  are both C,  $A^8$  is  $N-(CH_2)_d-R^7$  and  $A^9$  is N.

33. The use according to Claim 7 or 8, wherein  $R^1$  is an alkyl group, an O-alkyl group, F, Cl or Br,  $R^2$  and  $R^3$  are both H,  $X^1$  is NH, a is 1, b is 2,  $G^1$  is a group according to general formula 6,  $A^4$ ,  $A^5$  and  $A^{10}$  are all CH,  $A^6$  is NH,  $A^7$  and  $A^{11}$  are both C,  $A^8$  is  $N-(CH_2)_d-R^7$  and  $A^9$  is N.

34. The use according to Claim 7 or 8, wherein said compound is selected from the group consisting of:

- 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-(3,3-Dimethyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-(3-Methylsulfanyl-propyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-(2-Cyclopropyl-ethyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 3-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,

- 4-(2-Hydroxymethyl-cyclopropylmethyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-(3-Methyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 5 - 4-Cyclopentylmethyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclohexylmethyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 10 - 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 3-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 15 - 4-(2-Cyclopropyl-ethyl)-piperazine-1-carboxylic acid 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Pentyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 20 - 4-Hexyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- (R)-4-(2-Methyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-(2-Ethyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 25 - 4-(2-Methyl-but-2-enyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 30 - 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 2-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 2-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 35 4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,

- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 2-ethyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 5 - 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 2-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 2-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- and
- 10 - 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-methoxy-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide.

35. The use of a compound selected from the group consisting of:


- 4-(3,3-Dimethyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 15 - 4-(2-Cyclopropyl-ethyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 20 - 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 3-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-(2-Hydroxymethyl-cyclopropylmethyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 25 - 4-(3-Methyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclopentylmethyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclohexylmethyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 30 - 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 3-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 35 - 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,

- 4-(2-Cyclopropyl-ethyl)-piperazine-1-carboxylic acid 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 4-Pentyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 5 - 4-Hexyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - (R)-4-(2-Methyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 4-(2-Ethyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-
  - 10 dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 4-(2-Methyl-but-2-enyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 15 - 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 2-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 2-fluoro-4-(3-methyl-
  - 20 4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 2-ethyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 25 - 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 2-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 2-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - and
  - 30 - 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-methoxy-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- for the manufacture of a pharmaceutical composition for treatment of male erectile dysfunction.

- 35           36. A method for treatment of a disorder selected from the group consisting of primary dysmenorrhoea, pre-term labour, hypertension, Raynaud's

88


disease, brain oedema, motion sickness, small cell lung cancer, depression, anxiety, hyponatremia, liver cirrhosis and congestive heart failure which comprises the administration to a person in need of such treatment of therapeutically effective amount of a compound according to general formula 1, or a pharmaceutically acceptable salt thereof




1

wherein:


-  $G^1$  is selected from a group according to general formula 2, a group according to general formula 3, a group according to general formula 4, a group according to general formula 5, a group according to general formula 6 and a group according to general formula 7;




2




3




4



5



6



7

- A<sup>1</sup> is selected from CH<sub>2</sub>, CH(OH), NH, N-alkyl, O and S;
- A<sup>2</sup> is selected from CH<sub>2</sub>, CH(OH), C(=O) and NH;
- A<sup>3</sup> is selected from S, NH, N-alkyl, -CH=CH- and -CH=N-;
- A<sup>4</sup> and A<sup>5</sup> are each selected from CH and N;
- 5 - A<sup>6</sup> is selected from CH<sub>2</sub>, NH, N-alkyl and O;
- A<sup>7</sup> and A<sup>11</sup> are selected from C and N;
- A<sup>8</sup> and A<sup>9</sup> are selected from CH, N, NH, N(CH<sub>2</sub>)<sub>d</sub>R<sup>7</sup> and S;
- A<sup>10</sup> is selected from -CH=CH-, CH, N, NH, N(CH<sub>2</sub>)<sub>d</sub>R<sup>7</sup> and S;
- A<sup>12</sup> and A<sup>13</sup> are selected from N and C;
- 10 - A<sup>14</sup>, A<sup>15</sup> and A<sup>16</sup> are selected from NH, N-CH<sub>3</sub>, S, N and CH;
- X<sup>1</sup> is selected from O and NH;
- R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> are each selected from H, alkyl, O-alkyl, F, Cl and Br;
- R<sup>4</sup> is selected from H, alkyl, alkenyl, alkynyl, optionally substituted phenyl, optionally substituted thienyl, optionally substituted furyl, optionally substituted pyridyl, optionally substituted pyrrolyl, optionally substituted pyrazolyl,
- 15 optionally substituted imidazolyl, optionally substituted oxazolyl, optionally substituted isoxazolyl, optionally substituted thiazolyl, optionally substituted isothiazolyl, -(CH<sub>2</sub>)<sub>e</sub>R<sup>8</sup>, -CH<sub>2</sub>-CH=CH-CH<sub>2</sub>-R<sup>8</sup>, -CH<sub>2</sub>-C≡C-CH<sub>2</sub>-R<sup>8</sup>, -(CH<sub>2</sub>)<sub>g</sub>-CH(OH)-(CH<sub>2</sub>)<sub>h</sub>-R<sup>8</sup>, -(CH<sub>2</sub>)<sub>i</sub>-O-(CH<sub>2</sub>)<sub>j</sub>-R<sup>8</sup> and  $\text{CH}_2-\triangle-\text{R}^8$  ;
- 20 - R<sup>5</sup> and R<sup>6</sup> are independently selected from alkyl, Ar and -(CH<sub>2</sub>)<sub>f</sub>-Ar;
- R<sup>7</sup> is selected from H, alkyl, optionally substituted phenyl, F, OH, O-alkyl, O-acyl, S-alkyl, NH<sub>2</sub>, NH-alkyl, N(alkyl)<sub>2</sub>, NH-acyl, N(alkyl)-acyl, CO<sub>2</sub>H, CO<sub>2</sub>-alkyl, CONH<sub>2</sub>, CONH-alkyl, CON(alkyl)<sub>2</sub>, CN and CF<sub>3</sub>;
- R<sup>8</sup> is selected from H, alkyl, alkenyl, alkynyl, acyl, optionally substituted phenyl, optionally substituted pyridyl, optionally substituted thienyl, optionally substituted furyl, optionally substituted pyrrolyl, optionally substituted pyrazolyl, optionally substituted imidazolyl, optionally substituted oxazolyl, optionally substituted isoxazolyl, optionally substituted thiazolyl, optionally substituted isothiazolyl, F, OH, hydroxyalkyl, O-alkyl, O-acyl, S-alkyl, NH<sub>2</sub>,
- 25 NH-alkyl, N(alkyl)<sub>2</sub>, 1-pyrrolidinyl, 1-piperidinyl, 4-morpholinyl, NH-acyl, N(alkyl)-acyl, N<sub>3</sub>, CO<sub>2</sub>H, CO<sub>2</sub>-alkyl, CONH<sub>2</sub>, CONH-alkyl, CON(alkyl)<sub>2</sub>, CN and CF<sub>3</sub>;
- Ar is selected from optionally substituted thienyl and optionally substituted phenyl;

- a is 1 or 2, b is 1, 2 or 3; c is 1 or 2, d is 1, 2 or 3; e is 1, 2, 3 or 4; f is 1, 2 or 3 and g, h, i and j are all independently 1 or 2;

provided that:

- not more than one of  $A^8$ ,  $A^9$  and  $A^{10}$  is NH,  $N(CH_2)_dR^7$  or S;
- 5 -  $A^7$  and  $A^{11}$  are not both simultaneously N;
- neither  $A^7$  nor  $A^{11}$  is N if one of  $A^8$ ,  $A^9$  and  $A^{10}$  is NH,  $N(CH_2)_dR^7$  or S;
- if  $A^{10}$  is not  $-CH=CH-$  then one of  $A^8$ ,  $A^9$  and  $A^{10}$  is NH,  $N(CH_2)_dR^7$  or S or one of  $A^7$  and  $A^{11}$  is N;
- not more than one of  $A^{14}$ ,  $A^{15}$  and  $A^{16}$  is NH,  $N-CH_3$  or S;
- 10 -  $A^{12}$  and  $A^{13}$  are not both simultaneously N;
- if one of  $A^{14}$ ,  $A^{15}$  and  $A^{16}$  is NH,  $N-CH_3$  or S then  $A^{12}$  and  $A^{13}$  are both C;
- and
- one of  $A^{14}$ ,  $A^{15}$  and  $A^{16}$  is NH,  $N-CH_3$  or S or one of  $A^{12}$  and  $A^{13}$  is N.

15 37. The method of Claim 36, wherein at least one of  $R^1$ ,  $R^2$  and  $R^3$  is H and at least one is not H.

38. The method of Claim 36, wherein one of  $R^1$ ,  $R^2$  and  $R^3$  is selected from an alkyl group, an O-alkyl group, F, Cl and Br and the others are H.

20

39. The method of Claim 36, wherein  $X^1$  is NH.

40. The method of Claim 36, wherein a is 1 and b is 2.

25 41. The method of Claim 36, wherein  $G^1$  is a group according to general formula 3.

42. The method of to Claim 41, wherein c is 2.

30 43. The method of Claim 41, wherein  $A^1$  is  $CH_2$  and  $A^2$  is NH.

44. The method of Claim 41, wherein  $A^1$  is NH or N-alkyl and  $A^2$  is  $C(=O)$ .

35 45. The method of Claim 41, wherein  $A^3$  is S and  $A^4$  and  $A^5$  are both CH.



46. The method of Claim 41, wherein  $A^3$  is  $-\text{CH}=\text{CH}-$  and  $A^4$  and  $A^5$  are both CH.

5        47. The method of Claim 41, wherein  $A^3$  is  $-\text{CH}=\text{N}-$  and  $A^4$  and  $A^5$  are both CH.

10        48. The method of Claim 41, wherein  $A^3$  is  $-\text{CH}=\text{CH}-$ ,  $A^4$  is CH and  $A^5$  is N.

49. The method of Claim 36, wherein  $G^1$  is a group according to general formula 6 or 7

15        50. The method of Claim 49, wherein  $A^3$  is S and  $A^4$  and  $A^5$  are both CH.

51. The method of Claim 49, wherein  $A^3$  is  $-\text{CH}=\text{CH}-$  and  $A^4$  and  $A^5$  are both CH.

20        52. The method of Claim 49, wherein  $A^3$  is  $-\text{CH}=\text{N}-$  and  $A^4$  and  $A^5$  are both CH.

25        53. The method of Claim 49, wherein  $A^3$  is  $-\text{CH}=\text{CH}-$ ,  $A^4$  is CH and  $A^5$  is N.

54. The method of Claim 36, wherein  $G^1$  is a group according to general formula 4 or 6.

30        55. The method of Claim 54, wherein  $A^6$  is NH.

56. The method of Claim 54, wherein  $A^8$  is NH or  $\text{N}-(\text{CH}_2)_d-\text{R}^7$ .

57. The method of Claim 56, wherein  $A^9$  is N and  $A^{10}$  is CH.

58. The method of Claim 36, wherein one of  $R^1$ ,  $R^2$  and  $R^3$  is selected from an alkyl group, an O-alkyl group, F, Cl and Br and the others are H and  $X^1$  is NH.

5 59. The method of Claim 36, wherein one of  $R^1$ ,  $R^2$  and  $R^3$  is selected from an alkyl group, an O-alkyl group, F, Cl and Br and the others are H and  $X^1$  is NH, a is 1 and b is 2.

10 60. The method of Claim 36, wherein  $G^1$  is a group according to general formula 6,  $A^4$ ,  $A^5$  and  $A^{10}$  are all CH,  $A^6$  is NH,  $A^7$  and  $A^{11}$  are both C,  $A^8$  is N-(CH<sub>2</sub>)<sub>d</sub>- $R^7$  and  $A^9$  is N.

15 61. The method of Claim 36, wherein  $R^1$  is an alkyl group, an O-alkyl group, F, Cl or Br,  $R^2$  and  $R^3$  are both H,  $X^1$  is NH, a is 1, b is 2,  $G^1$  is a group according to general formula 6,  $A^4$ ,  $A^5$  and  $A^{10}$  are all CH,  $A^6$  is NH,  $A^7$  and  $A^{11}$  are both C,  $A^8$  is N-(CH<sub>2</sub>)<sub>d</sub>- $R^7$  and  $A^9$  is N.

20 62. The method of Claim 36, wherein said compound is selected from the group consisting of:  
- 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,  
- 4-(3,3-Dimethyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,  
- 4-(3-Methylsulfanyl-propyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,  
25 - 4-(2-Cyclopropyl-ethyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,  
- 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,  
30 - 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 3-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,  
- 4-(2-Hydroxymethyl-cyclopropylmethyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,  
35

- 4-(3-Methyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclopentylmethyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 5 - 4-Cyclohexylmethyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 3-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 10 - 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-(2-Cyclopropyl-ethyl)-piperazine-1-carboxylic acid 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 15 - 4-Pentyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Hexyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- (R)-4-(2-Methyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 20 - 4-(2-Ethyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-(2-Methyl-but-2-enyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 25 - 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 2-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 30 - 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 2-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 35 - 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 2-ethyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,

- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 2-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 2-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 5 and
- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-methoxy-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide.

63. A method for treatment of male erectile dysfunction which comprises the administration to a person in need of such treatment of therapeutically effective amount of a compound selected from the group consisting of:
- 4-(3,3-Dimethyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 4-(2-Cyclopropyl-ethyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 15 - 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 3-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 20 - 4-(2-Hydroxymethyl-cyclopropylmethyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 4-(3-Methyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 25 - 4-Cyclopentylmethyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 4-Cyclohexylmethyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 3-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 30 - 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
  - 35 - 4-(2-Cyclopropyl-ethyl)-piperazine-1-carboxylic acid 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,

- 4-Pentyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Hexyl-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 5 - (R)-4-(2-Methyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-(2-Ethyl-butyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-(2-Methyl-but-2-enyl)-piperazine-1-carboxylic acid 2-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 10 - 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-methyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 15 - 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 2-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 2-fluoro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 20 - 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 2-ethyl-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 2-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 25 - 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 2-chloro-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide,
- 4-Cyclobutylmethyl-piperazine-1-carboxylic acid 3-methoxy-4-(3-methyl-4,10-dihydro-3H-2,3,4,9-tetraaza-benzo[f]azulene-9-carbonyl)-benzylamide;
- and
- 30 pharmaceutically acceptable salts of the above mentioned compounds.

# INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/EP2004/001304

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D495/04 C07D487/04 C07D471/14 C07D471/04 C07D498/04  
A61K31/5517 A61P15/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data, EMBASE

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 03/016316 A (HUDSON PETER JEREMY ; ROE MICHAEL BRYAN (GB); ROOKER DAVID PHILIP ( ) 27 February 2003 (2003-02-27) page 3, line 1 - page 4, line 18; examples 54-56, 58, 60	1-63
A	WO 03/000692 A (ROE MICHAEL BRYAN ; ROOKER DAVID PHILIP (GB); PITT GARY ROBERT WILL) 3 January 2003 (2003-01-03) page 3, paragraph 1 - page 4, paragraph 2	1-63
A	US 5 760 031 A (ALBRIGHT JAY DONALD ET AL) 2 June 1998 (1998-06-02) column 1, line 10 - column 6, line 35	1-63
A	WO 02/083682 A (WYETH) 24 October 2002 (2002-10-24) page 7, line 1 - page 12, line 28 page 19, line 25 - page 20, line 5	1-63

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

9 June 2004

Date of mailing of the international search report

18/06/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Usueli, A

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2004/001304

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 03016316	A	27-02-2003	CA 2453962 A1	27-02-2003
			EP 1421087 A1	26-05-2004
			WO 03016316 A1	27-02-2003
WO 03000692	A	03-01-2003	CA 2450480 A1	03-01-2003
			EP 1399436 A2	24-03-2004
			WO 03000692 A2	03-01-2003
US 5760031	A	02-06-1998	US 5512563 A	30-04-1996
			US 5739128 A	14-04-1998
			US 5686445 A	11-11-1997
			US 5736538 A	07-04-1998
			US 5747487 A	05-05-1998
			US 5869483 A	09-02-1999
			US 5693635 A	02-12-1997
			US 5854236 A	29-12-1998
			US 5834461 A	10-11-1998
			US 5843952 A	01-12-1998
			US 5786353 A	28-07-1998
			AT 175198 T	15-01-1999
			AU 676737 B2	20-03-1997
			AU 6877694 A	09-02-1995
			BR 1100538 A3	08-08-2000
			CA 2128955 A1	30-01-1995
			CN 1106802 A ,B	16-08-1995
			CZ 9401798 A3	15-02-1995
			DE 69415614 D1	11-02-1999
			DE 69415614 T2	26-08-1999
			DK 640592 T3	30-08-1999
			EP 0640592 A1	01-03-1995
			ES 2125377 T3	01-03-1999
			FI 943542 A	30-01-1995
			FI 20011100 A	25-05-2001
			FI 20011101 A	25-05-2001
			FI 20011102 A	25-05-2001
			GR 3029758 T3	30-06-1999
			HK 1011362 A1	27-07-2001
			HU 71548 A2	28-12-1995
			JP 7179430 A	18-07-1995
			NO 942817 A	30-01-1995
			NZ 264116 A	26-11-1996
			NZ 299340 A	25-08-2000
			PL 304496 A1	06-02-1995
			RU 2149160 C1	20-05-2000
			SG 52312 A1	28-09-1998
			SK 88094 A3	10-05-1995
			TW 402592 B	21-08-2000
			ZA 9405604 A	09-03-1995
WO 02083682	A	24-10-2002	BR 0208943 A	27-04-2004
			CA 2443566 A1	24-10-2002
			EP 1377585 A1	07-01-2004
			WO 02083682 A1	24-10-2002
			US 2003004159 A1	02-01-2003